



## Intuitionistic Fuzzy Zweier I-convergent Double Sequence Spaces Defined by Orlicz Function

Vakeel A. Khan<sup>1</sup>, Yasmeen<sup>1</sup>, Hira Fatima<sup>1</sup>, Ayaz Ahmad<sup>2,\*</sup>

<sup>1</sup> Department of Mathematics, Aligarh Muslim University, Aligarh, India

<sup>2</sup> Department of Mathematics, National Institute of Technology, Patna, India

---

**Abstract.** The purpose of this paper is to introduce the intuitionistic fuzzy Zweier  $I$ -convergent double sequence spaces  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$  and  ${}_2\mathcal{Z}_{0(\mu,\nu)}^I(M)$  defined by Orlicz function and study the fuzzy topology on the said spaces.

**Key Words and Phrases:** Ideal, filter, double  $I$ -convergence, intuitionistic fuzzy normed spaces.

---

### 1. Introduction and Preliminaries

After the pioneering work of Zadeh [29], a huge number of research papers have been appeared on fuzzy theory and its applications as well as fuzzy analogues of the classical theories. Fuzzy set theory is a powerful hand set for modelling uncertainty and vagueness in various problems arising in field of science and engineering. It has a wide range of applications in various fields: population dynamics [3], chaos control [5], computer programming [6], nonlinear dynamical system [7], etc. Fuzzy topology is one of the most important and useful tools and it proves to be very useful for dealing with such situations where the use of classical theories breaks down. The concept of intuitionistic fuzzy normed space [25] and of intuitionistic fuzzy 2-normed space [21] are the latest developments in fuzzy topology. Recently V. A. Khan and Yasmeen ([12], [13]) studied the intuitionistic fuzzy Zweier  $I$ -convergent sequence spaces defined by modulus function and Orlicz function.

The notion of statistical convergence is a very useful functional tool for studying the convergence problems of numerical problems/matrices (double sequences) through the concept of density. The notion of  $I$ -convergence, which is a generalization of statistical convergence [4], was introduced by Kostyrko, Salat and Wilczynski [14] by using the idea of  $I$  of subsets of the set of natural numbers  $\mathbb{N}$  and further studied in [22]. Recently, the notion of statistical convergence of double sequences  $x = (x_{ij})$  has been defined and studied by Mursaleen and Edely [20]; and for fuzzy numbers by Savaş and Mursaleen [26]. Quite

---

\*Corresponding author.

Email addresses: vakhanmaths@gmail.com (V. A. Khan), yasmeen9828@gmail.com (Yasmeen), hirafatima2014@gmail.com (H. Fatima), ayaz1970@gmail.com (A. Ahmad)

recently, Das et al. [15] studied the notion of  $I$  and  $I^*$ - convergence of double sequences in  $\mathbb{R}$ .

We recall some notations and basic definitions used in this paper.

**Definition 1.** Let  $I \subset 2^{\mathbb{N}}$  be a non-trivial ideal in  $\mathbb{N}$ . Then a sequence  $x = (x_k)$  is said to be  $I$ -convergent to a number  $L$  if, for every  $\epsilon > 0$ , the set  $\{k \in \mathbb{N} : |x_k - L| \geq \epsilon\} \in I$ .

**Definition 2.** Let  $I \subset 2^{\mathbb{N}}$  be a non-trivial ideal in  $\mathbb{N}$ . Then a sequence  $x = (x_k)$  is said to be  $I$ -Cauchy if, for each  $\epsilon > 0$ , there exists a number  $N = N(\epsilon)$  such that the set  $\{k \in \mathbb{N} : |x_k - x_N| \geq \epsilon\} \in I$ .

**Definition 3.** The five-tuple  $(X, \mu, \nu, *, \diamond)$  is said to be an intuitionistic fuzzy normed space (for short, IFNS) if  $X$  is a vector space,  $*$  is a continuous  $t$ -norm,  $\diamond$  is a continuous  $t$ -conorm and  $\mu, \nu$  are fuzzy sets on  $X \times (0, \infty)$  satisfying the following conditions for every  $x, y \in X$  and  $s, t > 0$  :

- (a)  $\mu(x, t) + \nu(x, t) \leq 1$ ,
- (b)  $\mu(x, t) > 0$ ,
- (c)  $\mu(x, t) = 1$  if and only if  $x = 0$ ,
- (d)  $\mu(\alpha x, t) = \mu(x, \frac{t}{|\alpha|})$  for each  $\alpha \neq 0$ ,
- (e)  $\mu(x, t) * \mu(y, s) \leq \mu(x + y, t + s)$ ,
- (f)  $\mu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$  is continuous,
- (g)  $\lim_{t \rightarrow \infty} \mu(x, t) = 1$  and  $\lim_{t \rightarrow 0} \mu(x, t) = 0$ ,
- (h)  $\nu(x, t) < 1$ ,
- (i)  $\nu(x, t) = 0$  if and only if  $x = 0$ ,
- (j)  $\nu(\alpha x, t) = \nu(x, \frac{t}{|\alpha|})$  for each  $\alpha \neq 0$ ,
- (k)  $\nu(x, t) \diamond \nu(y, s) \geq \nu(x + y, t + s)$ ,
- (l)  $\nu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$  is continuous,
- (m)  $\lim_{t \rightarrow \infty} \nu(x, t) = 0$  and  $\lim_{t \rightarrow 0} \nu(x, t) = 1$ .

In this case  $(\mu, \nu)$  is called an intuitionistic fuzzy norm.

**Definition 4.** Let  $(X, \mu, \nu, *, \diamond)$  be an IFNS. Then a sequence  $x = (x_k)$  is said to be convergent to  $L \in X$  with respect to the intuitionistic fuzzy norm  $(\mu, \nu)$  if, for every  $\epsilon > 0$  and  $t > 0$ , there exists  $k_0 \in \mathbb{N}$  such that  $\mu(x_k - L, t) > 1 - \epsilon$  and  $\nu(x_k - L, t) < \epsilon$  for all  $k \geq k_0$ . In this case we write  $(\mu, \nu) - \lim x = L$ .

**Definition 5.** Let  $(X, \mu, \nu, *, \diamond)$  be an IFNS. Then a sequence  $x = (x_k)$  is said to be a Cauchy sequence with respect to the intuitionistic fuzzy norm  $(\mu, \nu)$  if, for every  $\epsilon > 0$  and  $t > 0$ , there exists  $k_0 \in \mathbb{N}$  such that  $\mu(x_k - x_l, t) < \epsilon$  and  $\nu(x_k - x_l, t) < \epsilon$  for all  $k, l \geq k_0$ .

**Definition 6.** Let  $K$  be the subset of natural numbers  $\mathbb{N}$ . Then the asymptotic density of  $K$ , denoted by  $\delta(K)$ , is defined as  $\delta(K) = \lim_n \frac{1}{n} |\{k \leq n : k \in K\}|$ , where the vertical bars denotes the cardinality of the enclosed set.

A number sequence  $x = (x_k)$  is said to be statistically convergent to a number  $\ell$  if, for each  $\epsilon > 0$ , the set  $K(\epsilon) = \{k \leq n : |x_k - \ell| > \epsilon\}$  has asymptotic density zero, i.e.

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - \ell| > \epsilon\}| = 0.$$

In this case we write  $st - \lim x = \ell$ .

**Definition 7.** A number sequence  $x = (x_k)$  is said to be statistically Cauchy sequence if, for every  $\epsilon > 0$ , there exists a number  $N = N(\epsilon)$  such that

$$\lim_n \frac{1}{n} |\{j \leq n : |x_j - x_N| \geq \epsilon\}| = 0.$$

The concepts of statistical convergence and statistical Cauchy for double sequences in intuitionistic fuzzy normed spaces have been studied by Mursaleen and Mohiuddine[14].

**Definition 8.** Let  $I \subset 2^{\mathbb{N}}$  be a non trivial ideal and  $(X, \mu, \nu, *, \diamond)$  be an IFNS. A sequence  $x = (x_k)$  of elements of  $X$  is said to be  $I$ -convergent to  $L \in X$  with respect to the intuitionistic fuzzy norm  $(\mu, \nu)$  if for every  $\epsilon > 0$  and  $t > 0$ , the set

$$\{k \in \mathbb{N} : \mu(x_k - L, t) \geq 1 - \epsilon \text{ or } \nu(x_k - L, t) \leq \epsilon\} \in I.$$

In this case  $L$  is called the  $I$ -limit of the sequence  $(x_k)$  with respect to the intuitionistic fuzzy norm  $(\mu, \nu)$  and we write  $I_{(\mu, \nu)} - \lim x_k = L$ .

## 2. $I_2$ -Convergence in an IFNS

**Definition 9.** Let  $(X, \mu, \nu, *, \diamond)$  be an IFNS. Then, a double sequence  $x = (x_{ij})$  is said to be statistically convergent to  $L \in X$  with respect to the intuitionistic fuzzy norm  $(\mu, \nu)$  if, for every  $\epsilon > 0$  and  $t > 0$ ,

$$\delta(\{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(x_{ij} - L, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - L, t) \geq \epsilon\}) = 0.$$

or equivalently

$$\lim_{mn} \frac{1}{mn} |\{i \leq m, j \leq n, : \mu(x_{ij} - L, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - L, t) \geq \epsilon\}| = 0.$$

In this case we write  $st_{(\mu, \nu)}^2 - \lim x = L$ .

**Definition 9.2** Let  $(X, \mu, \nu, *, \diamond)$  be an IFNS. Then, a double sequence  $x = (x_{ij})$  is said to be statistically Cauchy with respect to the intuitionistic fuzzy norm  $(\mu, \nu)$  if, for every

$\epsilon > 0$  and  $t > 0$ , there exist  $N = N(\epsilon)$  and  $M = M(\epsilon)$  such that for all  $i, p \geq N$  and  $j, q \geq M$ ,

$$\delta(\{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(x_{ij} - x_{pq}, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - x_{pq}, t) \geq \epsilon\}) = 0.$$

**Definition 10.** Let  $I_2$  be a non trivial ideal of  $\mathbb{N} \times \mathbb{N}$  and  $(X, \mu, \nu, *, \diamond)$  be an intuitionistic fuzzy normed space. A double sequence  $x = (x_{ij})$  of elements of  $X$  is said to be  $I_2$  convergent to  $L \in X$  with respect to the intuitionistic fuzzy norm  $(\mu, \nu)$  if, for each  $\epsilon > 0$  and  $t > 0$ ,

$$\{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(x_{ij} - L, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - L, t) \geq \epsilon\} \in I_2.$$

In this case we write  $I_2^{(\mu, \nu)} - \lim x = L$ .

The approach of constructing new sequence spaces by means of the matrix domain of a particular limitation method have been recently employed by Altay, Başar, Mursaleen [1], Malkowsky [19] Ng and Lee [23], and Wang [28]. Şengönül [27] defined the sequence  $y = (y_i)$  which is frequently used as the  $Z^p$  transformation of the sequence  $x = (x_i)$  i.e,

$$y_i = px_i + (1 - p)x_{i-1}$$

where  $x_{-1} = 0, p \neq 1, 1 < p < \infty$  and  $Z^p$  denotes the matrix  $Z^p = (z_{ik})$  defined by

$$z_{ik} = \begin{cases} p, & \text{if } (i = k), \\ 1 - p, & (i - 1 = k); (i, k \in \mathbb{N}) \\ 0, & \text{otherwise .} \end{cases}$$

Analogous to Başar and Altay [2], Şengönül [27] introduced the Zweier sequence spaces  $\mathcal{Z}$  and  $\mathcal{Z}_0$  as follows

$$\begin{aligned} \mathcal{Z} &= \{x = (x_k) \in \omega : \mathcal{Z}^p x \in c\}; \\ \mathcal{Z}_0 &= \{x = (x_k) \in \omega : \mathcal{Z}^p x \in c_0\}. \end{aligned}$$

Khan, Ebadullah and Yasmeen [8] introduced the following classes of sequences:

$$\mathcal{Z}^I = \{(x_k) \in \omega : \exists L \in \mathbb{C} \text{ such that for a given } \epsilon > 0, \{k \in \mathbb{N} : |x'_k - L| \geq \epsilon\} \in I\};$$

$$\mathcal{Z}_0^I = \{(x_k) \in \omega : \text{for a given } \epsilon > 0; \{k \in \mathbb{N} : |x'_k| \geq \epsilon\} \in I\}$$

,

where  $(x'_k) = (Z^p x)$ .

Recently V.A. Khan and Yasmeen [13] introduced the following sequence spaces:

$$\mathcal{Z}^I_{(\mu,\nu)}(M) = \left\{ (x_k) \in \omega : \left\{ k \in \mathbb{N} : M\left(\frac{\mu(x'_k - L, t)}{\rho}\right) \leq 1 - \epsilon \text{ or } M\left(\frac{\nu(x'_k - L, t)}{\rho}\right) \geq \epsilon \right\} \in I \right\},$$

$$\mathcal{Z}^I_{0(\mu,\nu)}(M) = \left\{ (x_k) \in \omega : \left\{ k \in \mathbb{N} : M\left(\frac{\mu(x'_k, t)}{\rho}\right) \leq 1 - \epsilon \text{ or } M\left(\frac{\nu(x'_k, t)}{\rho}\right) \geq \epsilon \right\} \in I \right\}.$$

**In this article we introduce the intuitionistic Zweier I-convergent double sequence spaces defined by Orlicz function as follows:**

$${}_2\mathcal{Z}^I_{(\mu,\nu)}(M) = \left\{ (x_{ij}) \in {}_2\omega : \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x'_{ij} - L, t)}{\rho}\right) \leq 1 - \epsilon \text{ or } M\left(\frac{\nu(x'_{ij} - L, t)}{\rho}\right) \geq \epsilon \right\} \in I_2 \right\};$$

$${}_2\mathcal{Z}^I_{0(\mu,\nu)}(M) = \left\{ (x_{ij}) \in {}_2\omega : \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x'_{ij}, t)}{\rho}\right) \leq 1 - \epsilon \text{ or } M\left(\frac{\nu(x'_{ij}, t)}{\rho}\right) \geq \epsilon \right\} \in I_2 \right\}.$$

### 3. Main results

**Theorem 1.** *The spaces  ${}_2\mathcal{Z}^I_{(\mu,\nu)}(M)$  and  ${}_2\mathcal{Z}^I_{0(\mu,\nu)}(M)$  are linear spaces.*

*Proof.* We prove the result for  ${}_2\mathcal{Z}^I_{(\mu,\nu)}(M)$ . Similarly the result can be proved for  ${}_2\mathcal{Z}^I_{0(\mu,\nu)}(M)$ . Let  $(x'_{ij}), (y'_{ij}) \in {}_2\mathcal{Z}^I_{(\mu,\nu)}(M)$  and let  $\alpha, \beta$  be scalars. Then for a given  $\epsilon > 0$ , we have

$$A_1 = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x'_{ij} - L_1, \frac{t}{2|\alpha|})}{\rho_1}\right) \leq 1 - \epsilon \text{ or } M\left(\frac{\nu(x'_{ij} - L_1, \frac{t}{2|\alpha|})}{\rho_1}\right) \geq \epsilon \right\} \in I_2;$$

$$A_2 = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(y'_{ij} - L_2, \frac{t}{2|\beta|})}{\rho_2}\right) \leq 1 - \epsilon \text{ or } M\left(\frac{\nu(y'_{ij} - L_2, \frac{t}{2|\beta|})}{\rho_2}\right) \geq \epsilon \right\} \in I_2. \text{ Thus}$$

$$A_1^c = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x'_{ij} - L_1, \frac{t}{2|\alpha|})}{\rho_1}\right) > 1 - \epsilon \text{ or } M\left(\frac{\nu(x'_{ij} - L_1, \frac{t}{2|\alpha|})}{\rho_1}\right) < \epsilon \right\} \in \mathcal{F}(I_2);$$

$$A_2^c = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(y'_{ij} - L_2, \frac{t}{2|\beta|})}{\rho_2}\right) > 1 - \epsilon \text{ or } M\left(\frac{\nu(y'_{ij} - L_2, \frac{t}{2|\beta|})}{\rho_2}\right) < \epsilon \right\} \in \mathcal{F}(I_2).$$

Define the set  $A_3 = A_1 \cup A_2$ , so that  $A_3 \in I_2$ . It follows that  $A_3^c$  is a non-empty set in  $\mathcal{F}(I_2)$ .

We shall show that for  $\rho_3 = \max\{2|\alpha| \rho_1, 2|\beta| \rho_2\}$  and for each  $(x'_{ij}), (y'_{ij}) \in {}_2\mathcal{Z}^I_{(\mu,\nu)}(M)$ ,

$$A_3^c \subset \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu((\alpha x'_{ij} + \beta y'_{ij}) - (\alpha L_1 + \beta L_2), t)}{\rho_3}\right) > 1 - \epsilon \text{ or}$$

$$M\left(\frac{\nu((\alpha x'_{ij} + \beta y'_{ij}) - (\alpha L_1 + \beta L_2), t)}{\rho_3}\right) < \epsilon \right\}.$$

Let  $(m, n) \in A_3^c$ . In this case

$$M\left(\frac{\mu(x_{mn}^{//} - L_1, \frac{t}{2|\alpha|})}{\rho_3}\right) > 1 - \epsilon \text{ or } M\left(\frac{\nu(x_{mn}^{//} - L_1, \frac{t}{2|\alpha|})}{\rho_3}\right) < \epsilon$$

and

$$M\left(\frac{\mu(y_{mn}^{//} - L_2, \frac{t}{2|\beta|})}{\rho_3}\right) > 1 - \epsilon \text{ or } M\left(\frac{\nu(y_{mn}^{//} - L_2, \frac{t}{2|\beta|})}{\rho_3}\right) < \epsilon.$$

we have

$$\begin{aligned} & M\left(\frac{\mu((\alpha x_{mn}^{//} + \beta y_{mn}^{//}) - (\alpha L_1 + \beta L_2), t)}{\rho_3}\right) \\ & \geq M\left(\frac{\mu(\alpha x_{mn}^{//} - \alpha L_1, \frac{t}{2})}{\rho_3}\right) * M\left(\frac{\mu(\beta y_{mn}^{//} - \beta L_2, \frac{t}{2})}{\rho_3}\right) \\ & = M\left(\frac{\mu(x_{mn}^{//} - L_1, \frac{t}{2|\alpha|})}{\rho_3}\right) * M\left(\frac{\mu(y_{mn}^{//} - L_2, \frac{t}{2|\beta|})}{\rho_3}\right) \\ & > (1 - \epsilon) * (1 - \epsilon) = (1 - \epsilon) \end{aligned}$$

and

$$\begin{aligned} & M\left(\frac{\nu((\alpha x_{mn}^{//} + \beta y_{mn}^{//}) - (\alpha L_1 + \beta L_2), t)}{\rho_3}\right) \\ & \leq M\left(\frac{\nu(\alpha x_{mn}^{//} - \alpha L_1, \frac{t}{2})}{\rho_3}\right) \diamond M\left(\frac{\nu(\beta y_{mn}^{//} - \beta L_2, \frac{t}{2})}{\rho_3}\right) \\ & = M\left(\frac{\nu(x_{mn}^{//} - L_1, \frac{t}{2|\alpha|})}{\rho_3}\right) \diamond M\left(\frac{\nu(y_{mn}^{//} - L_2, \frac{t}{2|\beta|})}{\rho_3}\right) \\ & > \epsilon \diamond \epsilon = \epsilon. \end{aligned}$$

This implies that

$$\begin{aligned} A_3^c \subset \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu((\alpha x_{ij}^{//} + \beta y_{ij}^{//}) - (\alpha L_1 + \beta L_2), t)}{\rho_3}\right) > 1 - \epsilon \text{ or } \right. \\ \left. M\left(\frac{\nu((\alpha x_{ij}^{//} + \beta y_{ij}^{//}) - (\alpha L_1 + \beta L_2), t)}{\rho_3}\right) < \epsilon \right\}. \end{aligned}$$

Hence  ${}_2\mathcal{Z}_{(\mu, \nu)}^I(M)$  is a linear space.

**Theorem 2.** Every open ball  ${}_2B_{x^{//}}(r, t)(M)$  is an open set in  ${}_2\mathcal{Z}_{(\mu, \nu)}^I(M)$ .

*Proof.* Let  ${}_2B_{x^{//}}(r, t)(M)$  be an open ball with centre  $x^{//}$  and radius  $r$  with respect to  $t$ .

That is

$${}_2B_{x^{//}}(r, t)(M) = \{(i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x_{ij}^{//} - L, t)}{\rho}\right) \leq 1 - r \text{ or } M\left(\frac{\nu(x_{ij}^{//} - L, t)}{\rho}\right) \geq r\} \in I_2.$$

Let  $y^{//} \in {}_2B_{x^{//}}^c(r, t)(M)$ . Then

$$M\left(\frac{\mu(x^{//} - y^{//}, t)}{\rho}\right) > 1 - r \text{ and } M\left(\frac{\nu(x^{//} - y^{//}, t)}{\rho}\right) < r$$

Since  $M\left(\frac{\mu(x^{//}-y^{//},t)}{\rho}\right) > 1-r$ , there exists  $t_0 \in (0,1)$  such that

$$M\left(\frac{\mu(x^{//}-y^{//},t_0)}{\rho}\right) > 1-r \quad \text{and} \quad M\left(\frac{\nu(x^{//}-y^{//},t_0)}{\rho}\right) < r.$$

Putting  $r_0 = M\left(\frac{\mu(x^{//}-y^{//},t_0)}{\rho}\right)$ . We have  $r_0 > 1-r$ , there exists  $s \in (0,1)$  such that  $r_0 > 1-s > 1-r$ .

For  $r_0 > 1-s$ , we have  $r_1, r_2 \in (0,1)$  such that  $r_0 * r_1 > 1-s$  and  $(1-r_0) \diamond (1-r_2) \leq s$ . Putting  $r_3 = \max\{r_1, r_2\}$ , consider the ball  ${}_2B_{y^{//}}^c(1-r_3, t-t_0)(M)$ . We prove that  ${}_2B_{y^{//}}^c(1-r_3, t-t_0)(M) \subset {}_2B_{x^{//}}^c(r, t)(M)$ .

Let  $z^{//} \in {}_2B_{y^{//}}^c(1-r_3, t-t_0)(M)$ .

$$M\left(\frac{\mu(y^{//}-z^{//},t-t_0)}{\rho}\right) > r_3 \quad \text{and} \quad M\left(\frac{\nu(y^{//}-z^{//},t-t_0)}{\rho}\right) < r_3.$$

Therefore,

$$\begin{aligned} M\left(\frac{\mu(x^{//}-z^{//},t)}{\rho}\right) &\geq M\left(\frac{\mu(x^{//}-y^{//},t_0)}{\rho}\right) * M\left(\frac{\mu(y^{//}-z^{//},t-t_0)}{\rho}\right) \\ &\geq (r_0 * r_3) \geq (r_0 * r_1) \geq (1-s) > (1-r). \end{aligned}$$

and

$$\begin{aligned} M\left(\frac{\nu(x^{//}-z^{//},t)}{\rho}\right) &\leq M\left(\frac{\nu(x^{//}-y^{//},t_0)}{\rho}\right) \diamond M\left(\frac{\nu(y^{//}-z^{//},t-t_0)}{\rho}\right) \\ &\leq (1-r_0) \diamond (1-r_3) \leq (1-r_0) > (1-r_2) < s < r. \end{aligned}$$

Thus  $z^{//} \in {}_2B_{x^{//}}^c(r, t)(M)$  and hence  ${}_2B_{y^{//}}^c(1-r_3, t-t_0)(M) \subset {}_2B_{x^{//}}^c(r, t)(M)$ .

**Remark 1.**  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$  is IFNS.

Define

$${}_2\tau_{(\mu,\nu)}(M) = \{A \subset {}_2\mathcal{Z}_{(\mu,\nu)}^I(M) : \text{for each } x \in A \text{ there exists } t > 0 \text{ and } r \in (0,1) \text{ s. t. } {}_2B_{x^{//}}^c(r, t)(M) \subset A\}$$

Then  ${}_2\tau_{(\mu,\nu)}(M)$  is a topology on  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$ .

**Theorem 3.** The topology  ${}_2\tau_{(\mu,\nu)}(M)$  on  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$  is first countable.

*Proof.*  $\left\{{}_2B_{x^{//}}\left(\frac{1}{n}, \frac{1}{n}\right)(M) : n = 1, 2, 3, \dots\right\}$  is a local base at  $x^{//}$  the topology  ${}_2\tau_{(\mu,\nu)}(M)$  on  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$  is first countable.

**Theorem 4.**  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$  and  ${}_2\mathcal{Z}_{0(\mu,\nu)}^I(M)$  are Housdorff spaces.

*Proof.* We prove the result for  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$ . Similarly the result can be proved for  ${}_2\mathcal{Z}_{0(\mu,\nu)}^I(M)$ .

Let  $x^{//}, y^{//} \in {}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$  such that  $x^{//} \neq y^{//}$ .

Then  $0 < M\left(\frac{\mu(x^{//}-y^{//},t)}{\rho}\right) < 1$

and  $0 < M\left(\frac{\nu(x^{//}-y^{//},t)}{\rho}\right) < 1$ .

Putting  $r_1 = M\left(\frac{\mu(x^{//}-y^{//},t)}{\rho}\right)$  and  $r_2 = M\left(\frac{\nu(x^{//}-y^{//},t)}{\rho}\right)$  and  $r = \max\{r_1, 1 - r_2\}$ .

For each  $r_0 \in (r, 1)$ , there exists  $r_3$  and  $r_4$  such that

$$r_3 * r_4 \geq r_0 \quad \text{and} \quad (1 - r_3) \diamond (1 - r_4) \leq (1 - r_0).$$

Putting  $r_5 = \max\{r_3, 1 - r_4\}$  and consider the open balls  ${}_2B_{x^{//}}(1 - r_5, \frac{t}{2})(M)$  and  ${}_2B_{y^{//}}(1 - r_5, \frac{t}{2})(M)$ .

Then clearly  ${}_2B_{x^{//}}(1 - r_5, \frac{t}{2}) \cap {}_2B_{y^{//}}(1 - r_5, \frac{t}{2})(M) = \phi$ .

For if there exists  $z^{//} \in {}_2B_{x^{//}}(1 - r_5, \frac{t}{2}) \cap {}_2B_{y^{//}}(1 - r_5, \frac{t}{2})(M)$ , then

$$\begin{aligned} r_1 &= M\left(\frac{\mu(x^{//}-y^{//},t)}{\rho}\right) \geq M\left(\frac{\mu(x^{//}-z^{//},\frac{t}{2})}{\rho}\right) * M\left(\frac{\mu(z^{//}-y^{//},\frac{t}{2})}{\rho}\right) \\ &\geq r_5 * r_5 \geq r_3 * r_3 \geq r_0 > r_1. \end{aligned}$$

and

$$\begin{aligned} r_2 &= M\left(\frac{\nu(x^{//}-y^{//},t)}{\rho}\right) \leq M\left(\frac{\nu(x^{//}-z^{//},\frac{t}{2})}{\rho}\right) \diamond M\left(\frac{\nu(z^{//}-y^{//},\frac{t}{2})}{\rho}\right) \\ &\leq (1 - r_5) \diamond (1 - r_5) \leq (1 - r_4) \diamond (1 - r_4) \leq (1 - r_0) < r, \end{aligned}$$

which is a contradiction.

Hence  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$  is Housdorff.

**Theorem 5.**  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(m)$  is an IFNS.  ${}_2\tau_{(\mu,\nu)}(M)$  is a topology on  ${}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$ . Then a sequence  $(x_{ij}^{//}) \in {}_2\mathcal{Z}_{(\mu,\nu)}^I(M)$ ,  $x_{ij}^{//} \rightarrow x^{//}$  if and if  $\mu(x_{ij}^{//} - x^{//}, t)(M)$  and

$$M\left(\frac{\nu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) \rightarrow 0$$

as  $i \rightarrow \infty, j \rightarrow \infty$ .

*Proof.* Fix  $t_0 > 0$ . Suppose  $x_{ij}^{//} \rightarrow x^{//}$ . Then for  $r \in (0, 1)$ , there exists  $n_0 \in \mathbb{N}$  such that  $x_{ij}^{//} \in {}_2B_{x^{//}}(r, t)(M)$  for all  $i \geq n_0, j \geq n_0$ .

$${}_2B_{x^{//}}(r, t)(M) = \{(i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) \leq 1 - r \text{ or } M\left(\frac{\nu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) \geq r\} \in I_2$$



such that  ${}_2B_{x//}^c(r, t)(M) \in \mathcal{F}(I_2)$ .

Then

$$1 - M\left(\frac{\mu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) < r$$

and  $M\left(\frac{\nu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) \rightarrow 0$  as  $i \rightarrow \infty, j \rightarrow \infty$

Conversely, if for each  $t > 0, M\left(\frac{\mu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) \rightarrow 1$  and

$$M\left(\frac{\nu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) \rightarrow 0$$

as  $i \rightarrow \infty, j \rightarrow \infty$ , then for  $r \in (0, 1)$ , there exists  $n_0 \in \mathbb{N}$  such that  $1 - M\left(\frac{\mu(x_{ij}^{//} - x^{//}, t)}{\rho}\right) < r$  for all  $i \geq n_0, j \geq n_0$ .

Thus  $x_{ij}^{//} \in {}_2B_{x_{ij}}^c(r, t)(M)$  for all  $i \geq n_0, j_0 \geq n_0$  and hence  $x_{ij}^{//} \rightarrow x^{//}$ .

**Theorem 6.** A double sequence  $x = (x_{ij}^{//}) \in {}_2\mathcal{Z}_{(\mu, \nu)}^I(M)$ . I-converges if and only if for every  $\epsilon > 0$  and  $t > 0$  there exists a number  $M = M(x, \epsilon, t), N = N(x, \epsilon, t)$  such that

$$\{(M, N) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x_{MN}^{//} - L, \frac{t}{2})}{\rho}\right) > 1 - \epsilon \text{ or } M\left(\frac{\nu(x_{MN}^{//} - L, \frac{t}{2})}{\rho}\right) < \epsilon\} \in \mathcal{F}(I_2).$$

*Proof.* Suppose that  ${}_2I_{(\mu, \nu)} - \lim x = L$  and let  $\epsilon > 0$  and  $t > 0$ . For a given  $\epsilon > 0$  choose,  $s > 0$  such that  $(1 - \epsilon) * (1 - \epsilon) > 1 - s$  and  $\epsilon \diamond \epsilon < s$ . Then for each  $x \in {}_2\mathcal{Z}_{(\mu, \nu)}^I(M)$ ,

$$A_x(\epsilon, t)(M) = \{(i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x_{ij}^{//} - L, \frac{t}{2})}{\rho}\right) \leq 1 - \epsilon \text{ or } M\left(\frac{\nu(x_{ij}^{//} - L, \frac{t}{2})}{\rho}\right) \geq \epsilon\} \in I_2$$

which implies that

$$A_x^c(\epsilon, t)(M) = \{(i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x_{ij}^{//} - L, \frac{t}{2})}{\rho}\right) > 1 - \epsilon \text{ or } M\left(\frac{\nu(x_{ij}^{//} - L, \frac{t}{2})}{\rho}\right) < \epsilon\} \in \mathcal{F}(I_2).$$

Conversely let us choose  $N \in A_x^c(\epsilon, t)(M)$ . Then

$$M\left(\frac{\mu(x_N^{//} - L, \frac{t}{2})}{\rho}\right) > 1 - \epsilon \text{ or } M\left(\frac{\nu(x_N^{//} - L, \frac{t}{2})}{\rho}\right) < \epsilon.$$

Now we want to show that there exists a number  $N = N(x, \epsilon, t)$  such that

$$\{(i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x_{ij}^{//} - x_N^{//}, t)}{\rho}\right) \leq 1 - s \text{ or } M\left(\frac{\nu(x_{ij}^{//} - x_N^{//}, t)}{\rho}\right) \geq s\} \in I_2.$$

For this, define for each  $x \in {}_2\mathcal{Z}_{(\mu, \nu)}^I(M)$ .

$${}_2B_x(\epsilon, t)(M) = \{(i, j) \in \mathbb{N} \times \mathbb{N} : M\left(\frac{\mu(x_{ij}^{//} - x_N^{//}, t)}{\rho}\right) \leq 1 - s \text{ or } M\left(\frac{\nu(x_{ij}^{//} - x_N^{//}, t)}{\rho}\right) \geq s\} \in I_2.$$

Now we show that  ${}_2B_x(\epsilon, t)(M) \subset {}_2A_x(\epsilon, t)(M)$ .

Suppose that  ${}_2B_x(\epsilon, t)(M) \subset {}_2A_x(\epsilon, t)(M)$ .

Then there exists  $(m, n) \in {}_2B_x(\epsilon, t)(M)$  and  $(m, n) \notin {}_2A_x(\epsilon, t)(M)$ . Therefore we have

$$M\left(\frac{\mu(x_{ij}'' - x_N'', t)}{\rho}\right) < 1 - s$$

and  $M\left(\frac{\mu(x_{ij}'' - L, \frac{t}{2})}{\rho}\right) > 1 - \epsilon$ .

In particular  $M\left(\frac{\mu(x_N'' - L, \frac{t}{2})}{\rho}\right) > 1 - \epsilon$ .

Therefore we have

$$\begin{aligned} 1 - s &\geq M\left(\frac{\mu(x_{mn}'' - x_N'', t)}{\rho}\right) \geq M\left(\frac{\mu(x_{mn}'' - L, \frac{t}{2})}{\rho}\right) * M\left(\frac{\mu(x_N'' - L, \frac{t}{2})}{\rho}\right) \\ &\geq (1 - \epsilon) * (1 - \epsilon) > 1 - s, \end{aligned}$$

which is not possible.

On the other hand  $M\left(\frac{\nu(x_{ij}'' - x_N'', t)}{\rho}\right) \geq s$  and  $M\left(\frac{\nu(x_{ij}'' - L, \frac{t}{2})}{\rho}\right) < \epsilon$ .

In particular  $M\left(\frac{\nu(x_N'' - L, \frac{t}{2})}{\rho}\right) < \epsilon$ .

Therefore we have

$$s \leq M\left(\frac{\nu(x_{mn}'' - x_N'', t)}{\rho}\right) \leq M\left(\frac{\nu(x_{mn}'' - L, \frac{t}{2})}{\rho}\right) \diamond M\left(\frac{\nu(x_N'' - L, \frac{t}{2})}{\rho}\right) \leq \epsilon \diamond \epsilon < s,$$

which is not possible.

Hence  ${}_2B_x(\epsilon, t)(M) \subset {}_2A_x(\epsilon, t)(M)$ ,  ${}_2A_x(\epsilon, t)(M) \in I_2$ .

This implies that  ${}_2B_x(\epsilon, t)(M) \in I_2$ . Hence proved.

## Acknowledgements

The authors would like to record their gratitude to the reviewer for his careful reading and making some useful corrections which improved the presentation of the paper.

## References

- [1] B. Altay, F. Başar, and Mursaleen, On the Euler sequence space which include the spaces  $\ell_p$  and  $\ell_\infty$ , Inform. Sci., 76(10), (2006) pp. 1450-1462.
- [2] F. Başar, and B. Altay, On the spaces of sequences of  $p$ -bounded variation and related matrix mappings, Ukrainion math. J. 55(2003).
- [3] L. C. Barros, R. C. Bassanezi, P. A. Tonelli, Fuzzy modelling in population dynamics, Ecol. Model, 128(2000)27-33.

- [4] H. Fast, Sur la convergence statistique, Colloq. Math.,2(1951), 241-244.
- [5] A. L. Fradkov, R. J. Evans, Control of chaos: Methods of applications in engineering, Chaos, solution and Fractals (29)(2005),33-56.
- [6] R. Giles, A computer program for fuzzy reasoning, Fuzzy sets and systems (4)(1980),221-234.
- [7] L. Hong, J. Q. Sun, Bifurcations of fuzzy non-linear dynamical systems, Commun. Nonlinear Sci.Numer. Simul, (1)(2006), 1-12.
- [8] V. A. Khan, K. Ebadullah and Yasmeen, On Zweier  $I$ -convergent sequence spaces, Proyecciones Journal of Mathematics Vol.(3)(33)(2014),259-276.
- [9] V. A. Khan, K. Ebadullah and R.K.A. Rababah, Intuitionistic fuzzy zweier  $I$ -convergent sequence spaces, Functinal Analysis: Theory, methods and applications, Vol. (1)(2015), 1-7.
- [10] V. A. Khan, Yasmeen, On Paranorm Type Intuitionistic Fuzzy Zweier  $I$ -convergent Sequence Spaces, New Trends in Mathematical Sciences(Submitted).
- [11] V. A. Khan, Yasmeen, Intuitionistic Fuzzy Zweier  $I$ -convergent Double Sequence Spaces, New Trends in Mathematical Sciences(In press).
- [12] V. A. Khan, Yasmeen, Intuitionistic Fuzzy Zweier  $I$ -convergent Sequence Spaces defined by Modulus function, (submitted).
- [13] V. A. Khan, Yasmeen, Intuitionistic Fuzzy Zweier  $I$ -convergent Sequence Spaces defined by Orlicz function, (submitted).
- [14] P. Kostyrko, T. Salat and W. Wilczynski,  $I$ -convergence, Real Analysis Exchange (26)(2000), No. 2, 669-686.
- [15] P. Das, P. Kostyrko, W. Wilczynski, P. Malik,  $I$  and  $I^*$ - convergence of double sequences, Math. Slovaca (58)(2008), 605-620.
- [16] I.J. Maddox, Spaces of strongly summable sequences, Qurt. Math, vol. (18)(1967), 345-355.
- [17] I.J. Maddox, Elements of functional analysis, Cambridge univ. Press (1970).
- [18] A. Wilansky, Summability through functional analysis, North Holland Mathematics Studies, Oxford (1984).
- [19] E. Malkowsky, Recent results in the theory of matrix transformation in sequence spaces, Math. Vesnik,(49)(1997),187-196.
- [20] M. Mursaleen, Osama H. H. Edely, statistical convergence of double sequences, J. Math. Anal. Appl. (288)(2003), 223-231.

- [21] M. Mursaleen, Q.M.D. Lohni, Intuitionistic fuzzy 2-normed space and some related concepts, *Chaos, solution and Fractals* (42)(2009), 331-344.
- [22] A. Nabiev, S. Pehlivan, M. Gürdal, On  $I$ - Cauchy sequence, *Taiwanese J. Math.* (11)(2)(2007), 569-576.
- [23] P. N. Ng and P.Y. Lee, Ceaaoro sequence spaces of non-absolute type, *Comment. Math.Pracc. Math.*(20)(2)(1978), 429-433.
- [24] J. H. Park, Intuitionistic fuzzy matric space, *Chaos, solution and Fractals* (22)(2004), 1039-1046.
- [25] R. Saddati, J. H. Park, On the intuitionistic fuzzy topological spaces, *Chaos, solution and Fractals* (27)(2006), 331-344.
- [26] E. Savaş, M. Mursaleen, On statistical convergent double sequences of fuzzy numbers, *Inform. Sci.* 162(2004), 183-192.
- [27] M. Sengönül, On the Zweier sequence space, *Demonstratio Mathematica*, Vol. XL No. (40)(2007), 181-196.1.
- [28] C. S. Wang, On Nörlund sequence spaces, *Tamkang J. Math.* (9)(1978), 269-274.1.
- [29] L. A. Zadeh, Fuzzy sets, *Inform Control*, (8)(1965), 338-353. 1, 1.8.