



Locally Conformal Almost Cosymplectic Manifold of Φ -holomorphic Sectional Conharmonic Curvature Tensor

Habeeb M. Abood¹, Farah Hassan J. Al-Hussaini^{1,*}

¹ *Department of Mathematics, College of Education for Pure Sciences, University of Basrah, Basrah, Iraq*

Abstract. The aim of the present paper is to study the geometry of locally conformal almost cosymplectic manifold of Φ -holomorphic sectional conharmonic curvature tensor. In particular, the necessary and sufficient conditions that locally conformal almost cosymplectic manifold is a manifold of point constant Φ -holomorphic sectional conharmonic curvature tensor have been found. The relation between the mentioned manifold and the Einstein manifold is determined.

2010 Mathematics Subject Classifications: 53C55, 53B35

Key Words and Phrases: Locally conformal almost cosymplectic manifold, conharmonic curvature tensor, Φ -holomorphic sectional conharmonic curvature tensor, Einstein manifold.

1. Introduction

Sectional curvature provides a lot of information with regard to substance geometry of Riemannian manifolds. Manifolds with constant sectional curvature are a great source of study. Moreover, contact geometry plays important roles in Physics, optics, differential equations and phase spaces of a dynamical system. This stimulated the researchers to work in the domain of constancy holomorphic sectional curvatures of locally conformal almost cosymplectic manifold which is a motivating class of almost contact metric manifold..

The study of constant holomorphic sectional curvature of almost Hermitian manifolds was started by Tanno [19] in 1973. He obtained an algebraic characterization for an almost Hermitian manifold to constringe to a space of constant holomorphic sectional curvature, which he later extended for Sasakian manifold. In 1988, Kim [7] studied total spaces of constant Φ -holomorphic sectional curvature and in 1989, he studied [8] total spaces with flat contact Bochner curvature tensor for fibred Sasakian spaces with conformal fibres. In 1993, Takano [18] discuss fibred Sasakian spaces of constant Φ -holomorphic sectional

*Corresponding author.

DOI: <https://doi.org/10.29020/nybg.ejpam.v11i3.3261>

Email addresses: iraqsafwan2006@gmail.com (H. M. Abood),
farahalhussaini14@yahoo.com (F. H. J. Al-Hussaini)

and at the same time Nagaich [14] showed a generalized Tanno's results for indefinite almost Hermitian manifold. In 2009, Rani et al. [17] considered similar condition of [19] to another distinct class of almost contact manifold known as (ϵ) -Sasakian manifold. In 2012, Kirichenko and Kharitonova [12] studied the constancy of Φ -holomorphic sectional curvature of normal locally conformal almost cosymplectic Manifold.

2. Preliminaries

In this section, we will focus our efforts on the study of almost contact metric manifold. In particular, we dedicate our study on the construction of the class of locally conformal almost cosymplectic manifold in the G -adjoined structure space.

Definition 2.1. [1] Let M be $2n + 1$ dimensional smooth manifold , η be differential 1-form called a contact form, ξ be vector field called a characteristic, Φ be an endomorphism of the module of the vector fields $X(M)$ called a structure endomorphism, then the triple (η, ξ, Φ) is called an almost contact structure if the following conditions hold

- (i) $\eta(\xi) = 1$;
- (ii) $\Phi(\xi) = 0$;
- (iii) $\eta \circ \Phi = 0$;
- (iv) $\Phi^2 = -id + \eta \otimes \xi$.

Moreover, if there is a Riemannian metric $g = \langle \cdot, \cdot \rangle$ on M such that $\langle \Phi X, \Phi Y \rangle = \langle X, Y \rangle - \eta(X)\eta(Y)$, $X, Y \in X(M)$, then the tetrad of tensors (η, ξ, Φ, g) is called an almost contact metric structure. In this case the manifold M equipped with this structure is called an almost contact metric manifold.

Definition 2.2. [9] Let (M, η, Φ, g) be almost contact metric manifold (\mathcal{AC} -manifold). In the module $X(M)$ we can determine two complementary projections m, ℓ , where $m = \eta \otimes \xi$ and $\ell = -\Phi^2$; thus $X(M) = L \oplus \mathfrak{N}$, where $L = Im\Phi = ker\eta$ and $\mathfrak{N} = Imm = ker\Phi$, where ℓ and m are the projections onto the submodules L and \mathfrak{N} respectively.

Definition 2.3. [9] In the complexification module L^c of the module L define two endomorphisms σ and $\bar{\sigma}$ as $\sigma = \frac{1}{2}(id - \sqrt{-1}\Phi)$ and $\bar{\sigma} = -\frac{1}{2}(id + \sqrt{-1}\Phi)$. We can define two projections by the forms:

$$\Pi = \sigma \circ \ell = -\frac{1}{2}(\Phi^2 - \sqrt{-1}\Phi) \text{ and } \bar{\Pi} = \bar{\sigma} \circ \ell = \frac{1}{2}(\Phi^2 + \sqrt{-1}\Phi),$$

where $\sigma \circ \Phi = \Phi \circ \sigma = i\sigma$ and $\bar{\sigma} \circ \Phi = \Phi \circ \bar{\sigma} = -i\bar{\sigma}$. Therefore, If we denote $Im\Pi = D_{\Phi}^{\sqrt{-1}}$ and $Im\bar{\Pi} = D_{\Phi}^{-\sqrt{-1}}$, then

$$X^c(M) = D_{\Phi}^{\sqrt{-1}} \oplus D_{\Phi}^{-\sqrt{-1}} \oplus D_{\Phi}^0,$$

where $D_{\Phi}^{\sqrt{-1}}$, $D_{\Phi}^{-\sqrt{-1}}$ and D_{Φ}^0 are proper submodules.

Definition 2.4. [11] *At each point $p \in M^{2n+1}$, there is a frame in $T_p^c(M)$ of the form $(p, \varepsilon_0, \varepsilon_1, \dots, \varepsilon_n, \varepsilon_{\hat{1}}, \dots, \varepsilon_{\hat{n}})$, where $\varepsilon_a = \sqrt{2}\sigma_p(e_p)$, $\varepsilon_{\hat{a}} = \sqrt{2}\bar{\sigma}(e_p)$, $\hat{a} = a + n$, $\varepsilon_0 = \xi_p$, the mappings $\sigma_p : L_p \rightarrow D_{\Phi}^{\sqrt{-1}}$, $\bar{\sigma}_p : L_p \rightarrow D_{\Phi}^{-\sqrt{-1}}$ are isomorphism and anti-isomorphism respectively, and e_a are orthonormal bases of L_p . The frame $(p, \varepsilon_0, \varepsilon_1, \dots, \varepsilon_n, \varepsilon_{\hat{1}}, \dots, \varepsilon_{\hat{n}})$ is called an A -frame.*

Lemma 2.1. [13] *The matrices components of tensors Φ_p and g_p in A -frame have the following forms respectively:*

$$(\Phi_j^i) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \sqrt{-1}I_n & 0 \\ 0 & 0 & -\sqrt{-1}I_n \end{pmatrix}, \quad (g_{ij}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -I_n \\ 0 & I_n & 0 \end{pmatrix},$$

where I_n is the identity matrix of order n .

It is well known, that the set of such frames defines an G -structure on M with structure group $1 \times U(n)$, represented by matrix of the form $\begin{pmatrix} 1 & 0 & 0 \\ 0 & A & 0 \\ 0 & 0 & A \end{pmatrix}$, where $A \in U(n)$. This structure is called an G -adjoined structure.

Definition 2.5. [1] *A skew-symmetric tensor $\Omega(X, Y) = g(X, \Phi Y)$ is called a fundamental form of the \mathcal{AC} -structure.*

Definition 2.6. [4] *An almost contact metric structure $S = (\eta, \xi, \Phi, g)$ is called an almost cosymplectic structure (\mathcal{AC}_f -structure) if*

(i) $d\eta = 0$;

(ii) $d\Omega = 0$.

Definition 2.7. [15] *A conformal transformation of an \mathcal{AC} -structure $S = (\eta, \xi, \Phi, g)$ on a manifold is the passage from S to an \mathcal{AC} -structure $\tilde{S} = (\tilde{\eta}, \tilde{\xi}, \tilde{\Phi}, \tilde{g})$ such that*

$$\tilde{\eta} = e^{-\sigma}\eta, \quad \tilde{\xi} = e^{\sigma}\xi, \quad \tilde{\Phi} = \Phi, \quad \tilde{g} = e^{-2\sigma}g$$

where σ is the determining function of the conformal transformation. If $\sigma = \text{const}$, then the conformal transformation is said to be trivial.

Definition 2.8. [15] *An \mathcal{AC} -structure S on a manifold M is said to be locally conformal almost cosymplectic (\mathcal{LCAC}_f -structure) if the restriction of this structure to some neighborhood U of an arbitrary point $p \in M$ admits a conformal transformation of an almost cosymplectic structure. This transformation is called a locally conformal. A manifold M equipped with an \mathcal{LCAC}_f -structure is called an \mathcal{LCAC}_f -manifold.*

Lemma 2.2. [6] *In the G -adjoined structure space, the collection of the structure equations of \mathcal{LCAC}_f -manifold has the following forms:*

- (i) $d\omega^a = -\omega_b^a \wedge \omega^b + B_c^{ab} \omega^c \wedge \omega_b + B^{abc} \omega_b \wedge \omega_c + B_b^a \omega \wedge \omega^b + B^{ab} \omega \wedge \omega_b;$
- (ii) $d\omega_a = \omega_a^b \wedge \omega_b + B_{ab}^c \omega_c \wedge \omega^b + B_{abc} \omega^b \wedge \omega^c + B_a^b \omega \wedge \omega_b + B_{ab} \omega \wedge \omega^b;$
- (iii) $d\omega = C_b \omega \wedge \omega^b + C^b \omega \wedge \omega_b;$
- (iv) $d\omega_b^a = -\omega_c^a \wedge \omega_b^c + A_b^{acd} \omega_c \wedge \omega_d + A_{bcd}^a \omega^c \wedge \omega^d + A_{bd}^{ac} \omega^d \wedge \omega_c + A_{bc0}^a \omega \wedge \omega^c + A_b^{ac0} \omega \wedge \omega_c;$

where

- (i) $B^{[abc]} = B_{[abc]} = 0;$
- (ii) $B^{[ab]} = B_{[ab]} = 0;$
- (iii) $B_b^a = B_a^b = \sigma_0 \delta_a^b;$
- (iv) $C^{ab} = C_{ab} = 0;$
- (v) $B_c^{ab} = 2\sigma^{[a} \delta_c^{b]}, B_{ab}^c = 2\sigma_{[a} \delta_b^c];$
- (vi) $C^b = -\sigma^b, C_b = -\sigma_b;$
- (vii) $A_b^{acd} = 2\delta_b^{[c} \sigma^{a]d} - 2\delta_b^{[d} \sigma^{a]c} + B_b^{acd} - 2\sigma^a \delta_b^{[d} \sigma^{c]} - 2\sigma_e B^{ae[d} \delta_b^{c]} + 2\sigma_b B^{abc};$
- (viii) $A_b^{[acd]} = \sigma_e B^{e[da} \delta_b^{c]};$
- (ix) $A_{[bd]}^{ac} = -2\delta_{[b}^{[c} \sigma_{d]}^a] + 2\sigma^{[a} \delta_b^{e]} \sigma_{[e} \delta_d^c] - 2\sigma^{[a} \delta_d^{e]} \sigma_{[e} \delta_b^c] + \frac{1}{2} B^{aec} B_{ebd};$
- (x) $A_b^{ac0} = -2\delta_b^{[c} \sigma^{a]0} + D_b^{ac} - \delta_b^a \sigma_0^c - 2B^{aec} B_{eb} - \sigma^a \sigma_0 \delta_b^c + 2B^{ac} \sigma_b - B^{ae} \sigma_e \delta_b^c;$
- (xi) $A_b^{[ac]0} = \sigma_0^{[c} \delta_b^{a]} - \sigma_d \delta_b^{[a} B^{d]c} + \sigma_0 \sigma^{[c} \delta_b^{a]} + \frac{1}{2} B^{dca} B_{bd};$
- (xii) $B^{a[bcd]} = -B^{a[db} \sigma^{c]};$
- (xiii) $B^{abc0} = -2D^{a[bc]} - B^{adc} \sigma_0;$
- (xiv) $\sigma^{[cd]} = \sigma_b B^{bcd}.$

Here $B^{abc}, B_{abc}; B^a, B_a; B_b^a, B_a^b; C^{ab}, C_{ab}; C^b, C_b; A_b^{acd}, A_{acd}^b; A_{bd}^{ac}; A_b^{ac0}, A_{ac0}^b; B^{abci}, B_{abci}; D^{abi}, D_{abi}$ and σ_{ij} are smooth functions in the G -adjoined structure space.

The following lemma gives the expression for the nonzero components of Riemannian curvature tensor of \mathcal{LCA}_f -manifold in the G -adjoined structure space.

Lemma 2.3. [6] *In the G -adjoined structure space, the components of Riemannian curvature tensor of \mathcal{LCA}_f -manifold have the following forms:*

(i) $R_{bcd}^a = 2(A_{bcd}^a + 4\sigma^{[a} \delta_{[c}^h] B_{d]hb} - \sigma_0 B_{b[d} \delta_{c]}^a);$

- (ii) $R_{bcd}^a = 2(2\delta_{[c}^{[b}\sigma_{d]}^a] + 2B^{hab}B_{hdc} - \delta_{[c}^a\delta_{d]}^b\sigma_0^2);$
- (iii) $R_{bcd}^a = A_{bc}^{ad} + 4\sigma^{[a}\delta_c^{h]} \sigma_{[h}\delta_b^d] - 4B^{dah}B_{chb} + B^{ad}B_{bc} - \delta_c^a\delta_b^d\sigma_0^2;$
- (iv) $R_{bcd}^{\hat{a}} = 2(2B_{[c|ab|d]} - 2\sigma_{[a}B_{b]cd} + B_{a[c}B_{d]b});$
- (v) $R_{0cd}^a = 2(\sigma_{0[c}\delta_{d]}^a + B^{ab}B_{bcd} - 2\sigma^{[a}\delta_{[c}^{h]}B_{d]h});$
- (vi) $R_{b\hat{c}0}^a = A_b^{ac0} + \sigma_b B^{ac} - \delta_b^c\sigma_0\sigma^a;$
- (vii) $R_{b\hat{c}0}^{\hat{a}} = 2B_{cab0} + 2B_{cab}\sigma_0;$
- (viii) $R_{0b0}^a = -\delta_b^a\sigma_{00} - \delta_b^a\sigma_0^2 - B_{cb}B^{ac} - \sigma_b^a - \sigma^a\sigma_b + 2\sigma^{[a}\delta_b^{c]} \sigma_c;$
- (ix) $R_{0\hat{b}0}^a = 2\sigma_0 B^{ab} - D^{ab0} - \sigma^{ab} - \sigma^a\sigma^b + 2B^{bac}\sigma_c.$

and the other components are conjugate to the above components or can be obtained by the property of symmetry for R or equal to zero.

Definition 2.9. [3] A Ricci tensor is a tensor of type $(2,0)$ which is defined by

$$r_{ij} = -R_{ijk}^k$$

Lemma 2.4. In the G -adjoined structure space, the components of the Ricci tensor of \mathcal{LCCAC}_f -manifold are given by the following forms:

- (i) $r_{ab} = 2(-2A_{(ab)c}^c - 4(\sigma^{[c}\delta_{[b}^{h]}B_{c]ha} + \sigma^{[c}\delta_{[a}^{h]}B_{c]hb}) + \sigma_0 B_{a[c}\delta_{b]}^c + \sigma_0 B_{b[c}\delta_{a]}^c + 2\sigma_0 B_{ab} - D_{ab0} - \sigma_{ab} - \sigma_a\sigma_b + 2B_{bah}\sigma^h);$
- (ii) $r_{\hat{a}b} = -4(\delta_{[b}^{[a}\sigma_{c]}^c] - \sigma_{[c}\delta_{[h}^b]\sigma^{[h}\delta_c^a]} - \frac{1}{2}\sigma^{[a}\delta_b^{h]}\sigma_h + B^{hca}B_{hcb} + B^{bch}B_{cha}) + (B^{cb}B_{ac} - B_{hb}B^{ah}) + A_{ac}^{cb} - \delta_b^a\sigma_{00} - 2n\sigma_0^2 - \sigma_b^a - \sigma^a\sigma_b;$
- (iii) $r_{a0} = -A_{ac0}^c - \sigma^c B_{ac} + n\sigma_0\sigma_a + 2(\sigma_{0[c}\delta_{a]}^c + B^{cb}B_{bca} - 2\sigma^{[c}\delta_{[c}^{h]}B_{a]h});$
- (iv) $r_{oo} = -2n(\sigma_{00} + \sigma_0^2) - 2B_{hc}B^{ch} - 2(\sigma_c^c + \sigma^c\sigma_c) + 4\sigma^{[c}\delta_c^{h]}\sigma_h.$

and the other components can be found by taking the conjugate operator to the above components.

Proof. The above components can be obtained directly from the Definition 2.10 and Lemma 2.5.

Definition 2.10. An \mathcal{LCCAC}_f -manifold has Φ -invariant Ricci tensor, if $\Phi \circ r = r \circ \Phi$.

Lemma 2.5. *An $\mathcal{LCA}\mathcal{C}_f$ -manifold has Φ -invariant Ricci tensor if and only if, in the G -adjoined structure space, the following condition*

$$r_{\hat{b}}^{\hat{a}} = r_{ab} = 0$$

holds.

We conclude this section by remembering the main concept of our study which is a conharmonic curvature tensor.

Definition 2.11. [5] *Let M be an \mathcal{AC} -manifold of dimension $2n + 1$. A tensor T of type $(4, 0)$ which is invariant under conharmonic transformation and defined by the form:*

$$T_{ijkl} = R_{ijkl} - \frac{1}{2n-1}(r_{il}g_{jk} - r_{jl}g_{ik} + r_{jk}g_{il} - r_{ik}g_{jl})$$

is called a conharmonic tensor, where $T_{ijkl} = -T_{jikl} = -T_{ijlk} = T_{klij}$.

Theorem 2.1. *In the G -adjoined structure space, the components of conharmonic curvature tensor of $\mathcal{LCA}\mathcal{C}_f$ -manifold are given by the following forms:*

- (i) $T_{abcd} = 2(2B_{[c|ab|d]} - 2\sigma_{[a}B_{b]cd} + B_{a[c}B_{d]b});$
- (ii) $T_{\hat{a}bcd} = 2(A_{bcd}^a + 4\sigma^{[a}\delta_c^{h]}B_{d]hb} - \sigma_0B_{b[d}\delta_c^a]) - \frac{1}{2n-1}(r_{bc}\delta_d^a - r_{bd}\delta_c^a);$
- (iii) $T_{\hat{a}bc\hat{d}} = A_{bc}^{ad} + 4\sigma^{[a}\delta_c^{h]} \sigma_{[h}\delta_b^d] - 4B^{dah}B_{chb} + B^{ad}B_{bc} - \delta_c^a\delta_b^d\sigma_0^2 - \frac{1}{2n-1}(r_b^d\delta_c^a + r_c^a\delta_b^d);$
- (iv) $T_{\hat{a}\hat{b}cd} = 2(2\delta_{[c}^{[b}\sigma_{d]}^a] + 2B^{hab}B_{hdc} - \delta_c^a\delta_d^b\sigma_0^2) - \frac{4}{2n-1}(r_{[a}^{[d}\delta_{b]}^c]);$
- (v) $T_{\hat{a}0cd} = 2(\sigma_0\delta_c^a\delta_d^a + B^{ab}B_{bcd} - 2\sigma^{[a}\delta_c^{h]}B_{d]h}) + \frac{1}{2n-1}(r_{0d}\delta_c^a - r_{0c}\delta_d^a);$
- (vi) $T_{\hat{a}b\hat{c}0} = A_b^{ac0} + \sigma_bB^{ac} - \delta_b^c\sigma_0\sigma^a - \frac{1}{2n-1}(r_0^a\delta_b^c);$
- (vii) $T_{abc0} = 2B_{cab0} + 2B_{cab}\sigma_0;$
- (viii) $T_{\hat{a}0b0} = -\delta_b^a\sigma_{00} - \delta_b^a\sigma_0^2 - B_{cb}B^{ac} - \sigma_b^a - \sigma^a\sigma_b + 2\sigma^{[a}\delta_b^{c]} \sigma_c + \frac{1}{2n-1}(r_{00}\delta_b^a + r_b^a);$
- (ix) $T_{\hat{a}0\hat{b}0} = 2\sigma_0B^{ab} - D^{ab0} - \sigma^{ab} - \sigma^a\sigma^b + 2B^{bac}\sigma_c + \frac{1}{2n-1}(r_{\hat{a}\hat{b}}).$

and the other components are conjugate to the above or can be obtained by the property of symmetry for T or equal to zero.

Definition 2.12. [16] *A Riemannian manifold is called an Einstein manifold, if the Ricci tensor satisfies the equation $r_{ij} = eg_{ij}$.*

Definition 2.13. [10] *Let M be an \mathcal{AC} -manifold, an Φ -holomorphic sectional curvature (Φ HS-curvature) of a manifold M in the direction $X \in X(M)$; $X \neq 0$ is a function $H(X)$ which is defined as:*

$$H(X) = \langle R(X, \Phi X, X, \Phi X), \rangle \|X\|^{-4}$$

Definition 2.14. [10] An \mathcal{AC} -manifold is called a manifold of point constant Φ HS-curvature if

$$\langle R(X, \Phi X, X, \Phi X,) \rangle = c\|X\|^4$$

where $c \in C^\infty(M)$; for all $X \in X(M)$

Lemma 2.6. [10] An \mathcal{AC} -manifold is a manifold of point constant Φ HS-curvature c if and only if, on the G -adjoined structure,

$$R^{(a}_{(bc)}{}^d) = \frac{c}{2}\delta_{bc}^{\bar{a}d}$$

where $\delta_{bc}^{\bar{a}d} = \delta_b^a\delta_c^d + \delta_c^a\delta_b^d$ is the symmetric second-order Kronecker delta.

Definition 2.15. Let M be an \mathcal{AC} -manifold, an Φ -holomorphic sectional conharmonic curvature (Φ HTS-curvature) of a manifold M in the direction $X \in X(M)$; $X \neq 0$ is a function $H(X)$ which is defined as

$$H(X) = \langle T(X, \Phi X, X, \Phi X,) \rangle \|X\|^{-4}$$

Definition 2.16. An \mathcal{AC} -manifold is called a manifold of point constant Φ HST-curvature if

$$\langle T(X, \Phi X, X, \Phi X,) \rangle = c\|X\|^4$$

where $c \in C^\infty(M)$; for all $X \in X(M)$.

3. The main results

This section is devoted to study the theoretical application of $\mathcal{LCA}\mathcal{C}_f$ -manifold of point constant Φ -holomorphic sectional conharmonic curvature. In particular, we found the necessary and sufficient conditions in which the $\mathcal{LCA}\mathcal{C}_f$ -manifold of point constant Φ -holomorphic sectional conharmonic curvature is an Eistein manifold.

The following theorems gives the necessary and sufficient condition in which an $\mathcal{LCA}\mathcal{C}_f$ -manifold is a manifold of point constant Φ HS-curvature.

Theorem 3.1. An $\mathcal{LCA}\mathcal{C}_f$ -manifold is a manifold of point constant Φ HS-curvature c if and only if, the relation $A^{(ad)}_{(bc)} = \frac{1}{2}\delta_{bc}^{\bar{a}d}(\sigma_0^2 + c) - 4\sigma^{[a}\delta_c^{h]}\sigma_{[h}\delta_b^d] + 4B^{(da)h}B_{chb} - B^{ad}B_{bc}$ holds on the G -adjoined structure space.

Proof. According to the components of the Riemannian curvature tensor of $\mathcal{LCA}\mathcal{C}_f$ -manifold, it follows that

$$R^a{}_{bc}{}^d = A^{ad}_{bc} + 4\sigma^{[a}\delta_c^{h]}\sigma_{[h}\delta_b^d] - 4B^{dah}B_{chb} + B^{ad}B_{bc} - \delta_c^a\delta_b^d\sigma_0^2$$

Symmetrizing with respect to the pair of upper and lower indices of the tensor $R^a{}_{bc}{}^d$, we get

$$R^{(a}{}_{(bc)}{}^d) = A^{(ad)}_{(bc)} + 4\sigma^{[a}\delta_c^{h]}\sigma_{[h}\delta_b^d] - 4B^{(da)h}B_{chb} + B^{ad}B_{bc} - \frac{1}{2}\delta_{bc}^{\bar{a}d}\sigma_0^2$$

By Lemma 2.6, the constancy condition on the Φ HST-curvature c for a \mathcal{LCA}_f -manifold, yields

$$A_{(bc)}^{(ad)} = \frac{1}{2}\delta_{bc}^{\hat{a}d}(\sigma_0^2 + c) - 4\sigma^{[a}\delta_c^{h]}\sigma_{[h}\delta_b^d] + 4B^{(da)h}B_{chb} - B^{ad}B_{bc}$$

Theorem 3.2. *Suppose that M is \mathcal{LCA}_f -manifold. Then the necessary and sufficient condition in which M is a manifold of point constant Φ HST-curvature C_0 is*

$$A_{bc}^{ad} = 4B^{dah}B_{chb} + B^{ad}B_{bc} - 4\sigma^{[a}\delta_c^{h]}\sigma_{[h}\delta_b^d] + \delta_c^a\delta_b^d\sigma_0^2 - C_0\delta_b^a\delta_c^d - \frac{1}{2n-1}(r_b^d\delta_c^a + r_c^a\delta_b^d)$$

Proof. Suppose that M is \mathcal{LCA}_f -manifold of the point constant Φ HST-curvature tensor.

According to the Definition 2.16, we get

$$\langle T(X, \Phi X, X, \Phi X,) \rangle = C_0\|X\|^4$$

In the G -adjoined structure space, we have

$$T_{ijkl}X^i(\Phi X)^jX^k(\Phi X)^l = C_0g_{ij}g_{kl}X^iX^jX^kX^l$$

According to the property $(\Phi X)^a = \sqrt{-1}X^a$, $(\Phi X)^{\hat{a}} = -\sqrt{-1}X^{\hat{a}}$ and $(\Phi X)^0 = 0$ and then using the properties of conharmonic tensor, we get

$$-4T_{\hat{a}bcd} = 4C_0\delta_b^a\delta_c^d$$

Hence

$$A_{bc}^{ad} = 4B^{dah}B_{chb} + B^{ad}B_{bc} - 4\sigma^{[a}\delta_c^{h]}\sigma_{[h}\delta_b^d] + \delta_c^a\delta_b^d\sigma_0^2 - C_0\delta_b^a\delta_c^d - \frac{1}{2n-1}(r_b^d\delta_c^a + r_c^a\delta_b^d)$$

Theorem 3.3. *If M is \mathcal{LCA}_f -manifold of point constant Φ HST-curvature tensor with flat holomorphic sectional curvature tensor and Φ -invariant Ricci tensor. Then M is an Einstein manifold.*

Proof. Suppose that M is a manifold of point constant Φ HST-curvature tensor.

According to Theorem 3.2, we have

$$A_{bc}^{ad} - 4B^{dah}B_{chb} - B^{ad}B_{bc} + 4\sigma^{[a}\delta_c^{h]}\sigma_{[h}\delta_b^d] + \frac{1}{2n-1}(r_b^d\delta_c^a + r_c^a\delta_b^d) = \delta_c^a\delta_b^d\sigma_0^2 - C_0\delta_b^a\delta_c^d \quad (3.1)$$

Symmetrizing and then antisymmetrizing (3.1) by the indices (a, h) and (a, d) respectively and since M is a manifold with flat holomorphic sectional curvature tensor, then we have

$$\frac{1}{2n-1}(r_b^{[d}\delta_c^{a]} + r_c^{[a}\delta_b^{d]}) = \frac{1}{2}(\delta_b^d\delta_c^a - \delta_c^d\delta_b^a)(\sigma_0^2 + C_0) \quad (3.2)$$

Contracting (3.2) by the indices (d, c) , we deduce

$$-\frac{(n-2)}{2(2n-1)}(r_b^a + r_d^d \delta_b^a) = -\frac{(n-1)}{2} \delta_b^a (\sigma_0^2 + C_0) \tag{3.3}$$

Symmetrizing and antisymmetrizing (3.3) by the indices (a, d) , we obtain

$$r_b^a = e \delta_b^a$$

where $e = \frac{(2n-1)(n-1)}{(n-2)}(\sigma_0^2 + C_0)$

Since the Ricci tensor is Φ -invariant

Therefore M is Einstein manifold.

Theorem 3.4. *If M is \mathcal{LCA}_f -manifold of point constant Φ HST-curvature tensor and Φ -invariant Ricci tensor, then M is an Einstein manifold if and only if $A_{bc}^{ac} = B^{ac}B_{bc} + c_1 \delta_b^a$*

Proof. Suppose that M is a manifold of point constant Φ HST-curvature tensor. According to the Theorem 3.2, we have

$$A_{bc}^{ad} = 4B^{dah}B_{chb} + B^{ad}B_{bc} - 4\sigma^{[a} \delta_c^{h]} \sigma_{[h} \delta_b^d] + \delta_c^a \delta_b^d \sigma_0^2 - C_0 \delta_b^a \delta_c^d - \frac{1}{(2n-1)}(r_b^d \delta_c^a + r_c^a \delta_b^d) \tag{3.4}$$

Symmetrizing (3.4) by the indices (a, h) , we get

$$A_{bc}^{ad} = B^{ad}B_{bc} + \delta_c^a \delta_b^d \sigma_0^2 - C_0 \delta_b^a \delta_c^d - \frac{1}{(2n-1)}(r_b^d \delta_c^a + r_c^a \delta_b^d) \tag{3.5}$$

Contracting (3.5) by the indices (c, d) , we deduce

$$A_{bc}^{ac} = B^{ac}B_{bc} + (\sigma_0^2 - nC_0)\delta_b^a - \frac{2}{(2n-1)}r_b^a \tag{3.6}$$

Since M is an Einstein manifold, it follows that

$$A_{bc}^{ac} = B^{ac}B_{bc} + c_1 \delta_b^a$$

where $c_1 = \sigma_0^2 - nC_0 - \frac{2e}{(2n-1)}$

Conversely, by substituted A_{bc}^{ac} in equation (3.6), we get

$$r_b^a = e \delta_b^a$$

According to Φ -invariant of Ricci tensor, it follows that M is Einstein manifold.

References

- [1] D.E. Blair, The theory of quasi-Sasakian structures, *J. Differential Geometry*, N. 1, P. 331-345, 1967.
- [2] D.E. Blair, *Riemannian Geometry of Contact and Symplectic Manifolds*, in *Progr. Math.* Birkhauser, Boston, MA, Vol. 203, 2002.
- [3] E. Cartan, *Riemannian Geometry in an Orthogonal Frame*, From lectures Delivered by E. Lie Cartan at the Sorbonne 1926-27, Izdat. Moskov. Univ., Moscow, 1960; World Sci., Singapore, 2001.
- [4] S.I. Goldberg and K. Yano, Integrability of almost cosymplectic structures, *Pacific Journal of Mathematics*, 31, 373-382, 1969.
- [5] Y. Ishi, On conharmonic transformation, *Tensor N. S.* 7, pp. 73-80, 1957.
- [6] S. V. Kharitonova, On the geometry of locally conformal almost cosymplectic manifolds, *Matematicheskie Zametki*, Vol. 86, No. 1, pp. 126-138, 2009.
- [7] B.H. Kim, Fibred Riemannian spaces with contact structure, *Hiroshima Math.J.*18, 493-508, 1988.
- [8] B. H. Kim, Fibred Sasakian spaces with vanishing contact Bochner curvature tensor, *ibid.* 19, 181-195, 1989.
- [9] V. F. Kirichenko, The method of generalization of Hermitian geometry in the almost Hermitian contact manifold, *Problems of geometry VINITI ANSSR*, V. 18, P. 25-71, 1986.
- [10] V. F. Kirichenko, Generalized quasi-Kaehlerian manifolds and axioms of CR-submanifolds in generalized Hermitian geometry I, *Geo. Dedicata* 51 (1), 75-104 (1994).
- [11] V. F. Kirichenko and A. R. Rustanov, *Differential Geometry of quasi Sasakian manifolds*, *Mathematical Collection*. 193(8): 71-100, 2002.
- [12] V. F. Kirichenko and S. V. Kharitonova, On the geometry of normal locally conformal almost cosymplectic manifolds, *Matematicheskie Zametki*, Vol. 91, No. 1, pp. 40-53, 2012.
- [13] V. F. Kirichenko, *Differential - Geometry structures on manifolds*, Second edition, expanded. Odessa, Printing House, 458 P., 2013.
- [14] R. K. Nagaich, Constancy of holomorphic sectional curvature in indefinite almost Hermitian manifolds, *KodaiMath.J.*16, 327, 1993.
- [15] Z. Olszak, Locally conformal almost cosymplectic manifolds, *Collq. Math.* 57(1), 73-87, 1989.

- [16] A. Z. Petrov, Einstein space, Phys-Math. Letr. Moscow, P. 463, 1961.
- [17] R. Rani, R. Kumar and R. K. Nagaich, Constancy of Φ -Holomorphic Sectional Curvature of (ϵ) -Sasakian Manifolds, ISSN 1995-0802, Lobachevskii Journal of Mathematics, Vol.30, No.1, pp.7680, 2009.
- [18] K. Takano, On Fibred Sasakian Space with Vanishing Contact Bochner Curvature Tensor, Colloquium Mathematicum, Vol. LXV, FASC.2, 1993.
- [19] S. Tanno, Constancy of holomorphic sectional curvature in almost Hermitian manifolds, KodaiMath. Sem. Rep.25, 190, 1973.