



On the aspects of enriched lattice-valued topological groups and closure of lattice-valued subgroups

T M G Ahsanullah^{1,*}, Fawzi Al-Thukair¹

¹ *Department of Mathematics, King Saud University, Riyadh, Saudi Arabia.*

Dedicated to Professor John N. Mordeson on the occasion of his 87th birthday

Abstract. Starting with \mathbb{L} as an enriched cl -premonoid, in this paper, we explore some categorical connections between \mathbb{L} -valued topological groups and Kent convergence groups, where it is shown that every \mathbb{L} -valued topological group determines a well-known Kent convergence group, and conversely, every Kent convergence group induces an \mathbb{L} -valued topological group. Considering an \mathbb{L} -valued subgroup of a group, we show that the category of \mathbb{L} -valued groups, $\mathbb{L}\text{-GRP}$ has initial structure. Furthermore, we consider a category $\mathbb{L}\text{-CLS}$ of \mathbb{L} -valued closure spaces, obtaining its relation with \mathbb{L} -valued Moore closure, and provide examples in relation to \mathbb{L} -valued subgroups that produce Moore collection. Here we look at a category of \mathbb{L} -valued closure groups, $\mathbb{L}\text{-CLGRP}$ proving that it is a topological category. Finally, we obtain a relationship between $\mathbb{L}\text{-GRP}$ and $\mathbb{L}\text{-TransTOLGRP}$, the category of \mathbb{L} -transitive tolerance groups besides adding some properties of \mathbb{L} -valued closures of \mathbb{L} -valued subgroups on \mathbb{L} -valued topological groups.

2020 Mathematics Subject Classifications: 03E72, 20N25, 18B05, 54A05, 54A20

Key Words and Phrases: Enriched lattice, \mathbb{L} -valued topology, \mathbb{L} -valued subgroup, \mathbb{L} -valued topological group, Moore collection, Moore closure, \mathbb{L} -valued closure group, Kent convergence group, category theory

1. Introduction

We have investigated a notion of \mathbb{L} -valued topological groups in [3], where we considered \mathbb{L} -valued subgroup of a group. Various aspects of \mathbb{L} -valued subgroups of groups are studied over the years by various authors, cf. [11, 23, 25, 26, 29] but its categorical behaviors are explored in a certain extent in recent times [26], although the category of fuzzy sets being studied for quite a long time, cf. [14, 33]. In [3], we also considered \mathbb{L} -valued closure of an \mathbb{L} -valued subgroup of a group in the context of \mathbb{L} -valued neighborhood groups, where the lattice under consideration was a complete MV-algebra with square roots.

Although our main objective of this paper is to explore further \mathbb{L} -valued subgroups from categorical view point and study category of \mathbb{L} -valued closure spaces vis-à-vis category of

*Corresponding author.

DOI: <https://doi.org/10.29020/nybg.ejpam.v14i3.4021>

Email addresses: tmga1@ksu.edu.sa (T.M.G. Ahsanullah), thukair@ksu.edu.sa (Fawzi Al-Thukair)

\mathbb{L} -valued closure groups in conjunction with \mathbb{L} -valued topological groups, we add some results on the connection of \mathbb{L} -valued topological groups and classical Kent convergence groups. However, we mainly focused on the impact of \mathbb{L} -valued closure structures on \mathbb{L} -valued topological groups instead of convergence groups. We arrange our work as follows. In Section 2, we give a short survey on \mathbb{L} -valued structures that we used in the text. The idea of convergence spaces and their connection to topological spaces is quite old, cf. [4–7, 10, 13, 20, 21, 27, 28]; following the concept of the compatibility of convergence structures with groups structures as proposed by D. C. Kent [20], for the first time, we explore a connection between the categories of \mathbb{L} -valued topological groups and Kent convergence groups, this is done in Section 3. We introduce the concept of \mathbb{L} -valued closure space, and \mathbb{L} -closure of \mathbb{L} -valued subgroup of a group in Section 4; we also introduce here a category of \mathbb{L} -valued closure groups - a topological category. With the help of connections, as presented by L. N. Stout in [32] and C. L. Waker in [33] between the categories of \mathbb{L} -**SET** and \mathbb{L} -**TOL**, the category of \mathbb{L} -valued tolerance spaces [32], we prove a connection between \mathbb{L} -**GRP**, category of \mathbb{L} -valued subgroups, and \mathbb{L} -valued transitive tolerance spaces, \mathbb{L} -**TranTOL**. Section 5 is devoted to study properties of \mathbb{L} -valued closure of \mathbb{L} -valued subgroups in the context of \mathbb{L} -valued topological groups, where some properties from groups are taken into consideration.

2. Preliminaries

Throughout the text we consider $\mathbb{L} = (\mathbb{L}, \leq)$ a complete lattice with \top , the top element and \perp , the bottom element of \mathbb{L} .

Definition 1. [16, 17] A triple $(\mathbb{L}, \leq, *)$, where $*$: $\mathbb{L} \times \mathbb{L} \longrightarrow \mathbb{L}$ is a binary operation on \mathbb{L} , is called a **GL-monoid** if and only if the following holds:

(GLM1) $(\mathbb{L}, *)$ is a commutative semigroup;

(GLM2) $\forall \alpha \in \mathbb{L}: \alpha * \top = \alpha$,

(GLM3) $*$ is distributive over arbitrary joins:

$\gamma * (\bigvee_{k \in K} \alpha_k) = \bigvee_{k \in K} (\gamma * \alpha_k)$, for $k \in K$, $\alpha_k, \gamma \in \mathbb{L}$;

(GLM4) for every $\gamma \leq \alpha$ there exists $\beta \in \mathbb{L}$ such that $\gamma = \alpha * \beta$ (divisibility).

The triple $(\mathbb{L}, \leq, *)$ is called a *commutative quantale* if (GLM1)-(GLM3) are fulfilled. If $*$ = \wedge , then the triple $(\mathbb{L}, \leq, \wedge)$ is called a frame or a complete Heyting algebra.

For a commutative quantale, the implication operator \rightarrow , also known as residuum, is given by

$$\rightarrow: \mathbb{L} \times \mathbb{L} \longrightarrow \mathbb{L}, \alpha \rightarrow \beta = \bigvee \{ \gamma \in \mathbb{L} \mid \alpha * \gamma \leq \beta \}.$$

A **GL-monoid** $(\mathbb{L}, \leq, *)$ is called a complete **MV-algebra** if

$$\forall \alpha \in \mathbb{L}, (\alpha \rightarrow \perp) \rightarrow \perp = \alpha \text{ (double negation).}$$

This means, in particular, that the unary operation $\neg: \mathbb{L} \longrightarrow \mathbb{L}, \alpha \mapsto \neg \alpha = \alpha \rightarrow \perp$ is an order-reversing involution.

Definition 2. [16, 17] A triple $(\mathbb{L}, \leq, \otimes)$, where $\otimes: \mathbb{L} \times \mathbb{L} \rightarrow \mathbb{L}$ is a binary operation on \mathbb{L} , is called a co-premonoid if and only if the following conditions are fulfilled:

- (CP1) $\forall \alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{L}: \alpha_1 \leq \beta_1$ and $\alpha_2 \leq \beta_2$ implies $\alpha_1 \otimes \alpha_2 \leq \beta_1 \otimes \beta_2$;
- (CP2) $\forall \alpha \in \mathbb{L}: \alpha \leq \alpha \otimes \top$ and $\alpha \leq \top \otimes \alpha$.

The category **COPML** consists of all co-premonoids as objects and morphisms as the mappings $\iota: (\mathbb{L}_1, \leq_1, \otimes_1) \rightarrow (\mathbb{L}_2, \leq_2, \otimes_2)$ satisfying the following conditions:

- (CPM1) ι preserves arbitrary joins;
- (CPM2) $\iota(\alpha \otimes_1 \alpha') = \iota(\alpha) \otimes_2 \iota(\alpha')$, $\forall \alpha, \alpha' \in \mathbb{L}_1$;
- (CPM3) ι preserves universal upper bounds; i.e., $\iota(\top) = \top$.

Definition 3. [16, 17] A co-premonoid $(\mathbb{L}, \leq, \otimes)$ is called a cl-premonoid if and only if (CP3) $\gamma \otimes (\bigvee_{k \in K} \alpha_k) = \bigvee_{k \in K} (\gamma \otimes \alpha_k)$, and $(\bigvee_{k \in K} \alpha_k) \otimes \gamma = \bigvee_{k \in K} (\alpha_k \otimes \gamma)$ for $K \neq \emptyset$, $k \in K$, $\alpha_k, \gamma \in \mathbb{L}$, is satisfied.

Definition 4. [16, 17] The quadruple $(\mathbb{L}, \leq, *, \otimes)$ is called an enriched cl-premonoid if and only if the following are fulfilled: (CLP1) $(\mathbb{L}, \leq, *)$ is a **GL**-monoid;

- (CLP2) $(\mathbb{L}, \leq, \otimes)$ is a cl-premonoid;
- (CLP3) $*$ is dominated by $\otimes: \forall \alpha, \beta, \gamma, \delta \in \mathbb{L}$,

$$(\alpha \otimes \beta) * (\gamma \otimes \delta) \leq (\alpha * \gamma) \otimes (\beta * \delta).$$

Definition 5. [16, 17] A **GL**-monoid $(\mathbb{L}, \leq, *)$ is said to have square roots if and only if there exists a unary operator $\mathcal{S}: \mathbb{L} \rightarrow \mathbb{L}$ such that the conditions below are satisfied:

- (S1) $\mathcal{S}(\alpha) * \mathcal{S}(\alpha) = \alpha$, $\forall \alpha \in \mathbb{L}$;
- (S2) $\beta * \beta \leq \alpha$ implies $\beta \leq \mathcal{S}(\alpha)$.

Since the formation of square roots is uniquely determined by (S1) and (S2), $\mathcal{S}(\alpha)$ is also written as $\alpha^{\frac{1}{2}}$.

A **GL**-monoid with square roots satisfies (S3) if it fulfills the following axiom:

- (S3) $(\alpha * \beta)^{\frac{1}{2}} = (\alpha^{\frac{1}{2}} * \beta^{\frac{1}{2}}) \vee \perp^{\frac{1}{2}}$, $\forall \alpha, \beta \in \mathbb{L}$.

If $\mathbb{L} = (\mathbb{L}, \leq, *)$ is a **GL**-monoid with square roots, then the monoidal mean operator $\otimes: \mathbb{L} \times \mathbb{L} \rightarrow \mathbb{L}$ is given by

$$\alpha \otimes \beta = \alpha^{\frac{1}{2}} * \beta^{\frac{1}{2}}, \forall \alpha, \beta \in \mathbb{L}.$$

An enriched cl-premonoid $\mathbb{L} = (\mathbb{L}, \leq, *, \otimes)$ is said to be pseudo-bisymmetric if it satisfies the following axiom:

$$(\alpha * \beta) \otimes (\gamma * \delta) = ((\alpha \otimes \gamma) * (\beta \otimes \delta)) \vee ((\alpha \otimes \perp) * (\beta \otimes \top)) \vee ((\perp \otimes \gamma) * (\top \otimes \delta)), \forall \alpha, \beta, \gamma, \delta \in \mathbb{L}.$$

Remark 1. [16, 17] (1) If $(L, \leq, *)$ is a **GL**-monoid with square roots, satisfying (S3), and \otimes is the monoidal mean operator \otimes , then the quadruple $(\mathbb{L}, \leq, *, \otimes)$ is pseudo-bisymmetric.

(2) If the cl-premonoid operation \otimes is identical to the quantal operation $*$, that is, $\otimes = *$, then the triple $(\mathbb{L}, \leq, *, \otimes)$ is pseudo-bisymmetric.

Proposition 1. [18] Let $(\mathbb{L}, \leq, *)$ be a *GL-monoid*. Then the following are fulfilled

$\forall \alpha, \beta, \gamma, \delta, \alpha_j, \beta_j, \gamma_j \in \mathbb{L}$:

- (1) $\alpha \leq \beta \rightarrow \gamma \Leftrightarrow \alpha * \beta \leq \gamma$;
- (2) $\alpha * (\alpha \rightarrow \beta) \leq \beta$;
- (3) $\alpha \leq \beta \Rightarrow \alpha \rightarrow \gamma \leq \beta \rightarrow \gamma$;
- (4) $\alpha \leq \beta \Rightarrow \gamma \rightarrow \alpha \geq \gamma \rightarrow \beta$;
- (5) $(\alpha \rightarrow \beta) \rightarrow \beta \geq \alpha$;
- (6) $\alpha * (\beta \rightarrow \gamma) \leq \beta \rightarrow (\alpha * \gamma)$;
- (7) $\alpha \rightarrow (\bigwedge_{j \in J} \beta_j) = \bigwedge_{j \in J} (\alpha \rightarrow \beta_j)$;
- (8) $(\bigvee_{j \in J} \alpha_j) \rightarrow \beta = \bigwedge_{j \in J} (\alpha_j \rightarrow \beta)$;
- (9) if $\alpha, \beta \in \mathbb{L}$ with $\alpha \leq \beta$, then for any $\gamma \in \mathbb{L}$, $\gamma * \alpha \leq \gamma * \beta$;
- (10) $\bigwedge_{j \in J} (\alpha_j * \gamma_j) \geq (\bigwedge_{j \in J} \alpha_j) * (\bigwedge_{j \in J} \gamma_j)$;
- (11) $(\alpha \rightarrow \gamma) * (\beta \rightarrow \delta) \leq \alpha * \beta \rightarrow \gamma * \delta$;
- (12) $\alpha \leq \beta \Leftrightarrow \alpha \rightarrow \beta = \top$;
- (13) $\alpha \rightarrow \top = \top$, $\top \rightarrow \alpha = \alpha$, and $\perp \rightarrow \alpha = \top$.

In what follows, the quadruple $\mathbb{L} = (\mathbb{L}, \leq, *, \otimes)$ (or simply \mathbb{L}) is assumed to be an enriched *cl-premonoid*, where $*$ is reserved for the *GL-monoid* operation, \otimes is for *cl-premonoid*, unless otherwise specified. The set of all \mathbb{L} -sets or \mathbb{L} -valued sets and is denoted by $\mathbb{L}^X (= \{\nu: X \rightarrow \mathbb{L}\})$. If $f: X \rightarrow Y$ is a function, then $f^{\leftarrow}: \mathbb{L}^Y \rightarrow \mathbb{L}^X$ is defined for any $\mu \in \mathbb{L}^Y$ by $f^{\leftarrow}(\mu) = \mu \circ f$; and $f^{\rightarrow}: \mathbb{L}^X \rightarrow \mathbb{L}^Y$ is defined by

$$f^{\rightarrow}(\nu)(y) = \bigvee \{\nu(x) \mid f(x) = y\},$$

for all $\nu \in \mathbb{L}^X, y \in Y$.

If \cdot is a binary operation on a set X , then we define the binary operation \odot on \mathbb{L}^X as follows. For $\nu_1, \nu_2 \in \mathbb{L}^X$ and $z \in X$

$$\nu_1 \odot \nu_2(z) = \bigvee \{\nu_1(x) * \nu_2(y) \mid x, y \in X, x \cdot y = z\};$$

usually, we write xy instead of $x \cdot y$. If $\nu_1, \nu_2 \in \mathbb{L}^X$, and $\rightarrow, *, \otimes$ are operations on \mathbb{L} as explained before, then these operations are carried over to \mathbb{L}^X point-wise:

- (i) $(\nu_1 \rightarrow \nu_2)(x) = \nu_1(x) \rightarrow \nu_2(x)$;
- (ii) $(\nu_1 * \nu_2)(x) = \nu_1(x) * \nu_2(x)$;
- (iii) $(\nu_1 \otimes \nu_2)(x) = \nu_1(x) \otimes \nu_2(x), \forall x \in X$.

Definition 6. [17, 18] A map $\mathcal{F}: \mathbb{L}^X \rightarrow \mathbb{L}$ is called an \mathbb{L} -valued filter on X if and only if the conditions below are satisfied:

- (LF1) $\mathcal{F}(\top_X) = \top, \mathcal{F}(\perp_X) = \perp$;
- (LF2) if $\nu_1, \nu_2 \in \mathbb{L}^X$ with $\nu_1 \leq \nu_2$, then $\mathcal{F}(\nu_1) \leq \mathcal{F}(\nu_2)$;
- (LF3) $\mathcal{F}(\nu_1) \otimes \mathcal{F}(\nu_2) \leq \mathcal{F}(\nu_1 \otimes \nu_2), \forall \nu_1, \nu_2 \in \mathbb{L}^X$.
- (S \mathbb{L}) An \mathbb{L} -valued filter \mathcal{F} is called a stratified \mathbb{L} -valued filter if $\forall \alpha \in \mathbb{L}, \forall \mu \in \mathbb{L}^X, \alpha * \mathcal{F}(\mu) \leq \mathcal{F}(\alpha * \mu)$.

The set of all stratified \mathbb{L} -valued filters on X is denoted by $\mathcal{F}_{\mathbb{L}}^s(X)$. On $\mathcal{F}_{\mathbb{L}}^s(X)$, partial ordering \leq is defined by: if $\mathcal{F}, \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(X)$, then $\mathcal{F} \leq \mathcal{G} \Leftrightarrow \mathcal{F}(\nu) \leq \mathcal{G}(\nu), \forall \nu \in \mathbb{L}^X$. If

$x \in X$, then $[x] \in \mathcal{F}_{\mathbb{L}}^s(X)$, called point stratified \mathbb{L} -valued filter on X , and is defined as $[x](\nu) = \nu(x)$, for all $\nu \in \mathbb{L}^X$.

If $\mathcal{F} \in \mathcal{F}_{\mathbb{L}}^s(X)$, then the stratified \mathbb{L} -valued filter $f^{\Rightarrow}(\mathcal{F}): \mathbb{L}^Y \rightarrow \mathbb{L}$ on Y is defined for any $\mu \in \mathbb{L}^Y$ by

$$[f^{\Rightarrow}(\mathcal{F})](\mu) = \mathcal{F}(f^{\Leftarrow}(\mu)) = \mathcal{F}(\mu \circ f).$$

If $\mathcal{F} \in \mathcal{F}_{\mathbb{L}}^s(Y)$, then $f^{\Leftarrow}(\mathcal{F}): \mathbb{L}^X \rightarrow \mathbb{L}$ is defined by

$$[f^{\Leftarrow}(\mathcal{F})](\nu) = \bigvee \{ \mathcal{F}(\mu) \mid \mu \in \mathbb{L}^Y, f^{\Leftarrow}(\mu) \leq \nu \},$$

for all $\nu \in \mathbb{L}^X$, is a stratified \mathbb{L} -filter on X if and only if for all $\mu \in \mathbb{L}^Y, f^{\Leftarrow}(\mu) = \perp_X \Rightarrow \mathcal{F}(\mu) = \perp$.

If $\nu \in \mathbb{L}^X$ and $\mu \in \mathbb{L}^Y$, then the product $\nu \times \mu: X \times Y \rightarrow \mathbb{L}$ is defined by:

$$\nu \times \mu = \nu \circ pr_1 * \mu \circ pr_2,$$

where $pr_1: X \times Y \rightarrow X, (x, y) \mapsto x$ and $pr_2: X \times Y \rightarrow Y, (x, y) \mapsto y$ are usual projections. Note that in the preceding definition of product \mathbb{L} -set the operation $*$ holds only for finite case; otherwise, we need to take $*$ = \wedge .

Proposition 2. [16] *If $(L, \leq, *)$ is a GL -monoid, then for stratified L -valued filters \mathcal{F}_1 and \mathcal{F}_2 , the supremum $\mathcal{F}_1 \vee \mathcal{F}_2$ exists if and only if $\mathcal{F}_1(\nu_1) * \mathcal{F}_2(\nu_2) = \perp \forall \nu_1, \nu_2 \in L^X$ such that $\nu_1 * \nu_2 = \perp_X$. In particular, the supremum is the stratified L -valued filter defined for all $\nu \in L^X$ by*

$$\mathcal{F}_1 \vee \mathcal{F}_2(\nu) = \bigvee \{ \mathcal{F}_1(\nu_1) * \mathcal{F}_2(\nu_2) \mid \nu_1, \nu_2 \in L^X, \nu_1 * \nu_2 \leq \nu \}.$$

Let (G, \cdot) be a group. If $\mathcal{F} \in \mathbb{L}^s(G)$, then \mathcal{F}^{-1} is defined by $\mathcal{F}^{-1}(\nu) = \mathcal{F}(\nu^{-1})$, where $\nu^{-1}: G \rightarrow \mathbb{L}, x \mapsto \nu(x^{-1})$. Clearly, $\mathcal{F}^{-1} \in \mathcal{F}_{\mathbb{L}}^s(G)$, since for any $\nu \in L^X, j^{\Rightarrow}(\mathcal{F})(\nu) = \mathcal{F}(j^{\Leftarrow}(\nu)) = \mathcal{F}(\nu^{-1}) = \mathcal{F}^{-1}(\nu)$, where $j: G \rightarrow G, x \mapsto x^{-1}$. Also, if $m: G \times G \rightarrow G, (g, h) \mapsto gh$, then for any $\nu_1, \nu_2 \in \mathbb{L}^G$ and $z \in G, m^{\Rightarrow}(\nu_1 \times \nu_2)(z) = \bigvee_{m(g,h)=z} (\nu_1 \times \nu_2)(g, h) = \bigvee_{gh=z} (\nu_1 \circ pr_1 * \nu_2 \circ pr_2)(g, h) = \bigvee_{gh=z} \nu_1 \circ pr_1(g, h) * \nu_2 \circ pr_2(g, h) = \bigvee_{gh=z} \nu_1(g) * \nu_2(h) = \nu_1 \odot \nu_2(z)$.

Lemma 1. [3] *Let $\mathbb{L} = (\mathbb{L}, \leq, *)$ be a GL -monoid and $(G, \cdot) \in |GRP|$. Then for any $\mathcal{F}, \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(X), m^{\Rightarrow}(\mathcal{F} \times \mathcal{G}) = \mathcal{F} \odot \mathcal{G}$.*

Definition 7. [17] *Consider a mapping $\mathfrak{N}: X \rightarrow \mathbb{L}^{\mathbb{L}^X}$ such that the following conditions are fulfilled:*

- (LN1) $\mathfrak{N}^x(\top_X) = \top$;
- (LN2) $\mathfrak{N}^x(\nu_1) \leq \mathfrak{N}^x(\nu_2)$ for all $\nu_1, \nu_2 \in \mathbb{L}^X$ with $\nu_1 \leq \nu_2$;
- (LN3) $\mathfrak{N}^x(\nu_1) \otimes \mathfrak{N}^x(\nu_2) \leq \mathfrak{N}^x(\nu_1 \otimes \nu_2)$, for all $\nu_1, \nu_2 \in \mathbb{L}^X$;
- (LN4) $\mathfrak{N}^x(\nu) \leq \nu(x)$, for all $\nu \in \mathbb{L}^X$;
- (LN5) $\forall x \in X$ and $\nu \in \mathbb{L}^X, \mathfrak{N}^x(\nu) \leq \bigvee \{ \mathfrak{N}^x(\mu) : \mu \in \mathbb{L}^X, \mu(y) \leq [\mathfrak{N}^y](\nu), \forall y \in X \}$
- (SLN) $\alpha * \mathfrak{N}^x(\nu) \leq \mathfrak{N}^x(\alpha * \nu)$.

Then $\mathfrak{N} = (\mathfrak{N}^x)_{x \in X}$ is called a stratified \mathbb{L} -valued neighborhood system on X , and the

pair $(X, \mathfrak{N} = (\mathfrak{N}^x)_{x \in X})$ is called a stratified \mathbb{L} -valued neighborhood space.

If (X, \mathfrak{N}) and (Y, \mathfrak{M}) stratified \mathbb{L} -valued neighborhood spaces, then a map $f: (X, \mathfrak{N}) \rightarrow (Y, \mathfrak{M})$ is said to be continuous at a point $x \in X$ if and only if $\mathfrak{M}^{f(x)}(\nu) \leq \mathfrak{N}^x(f^{\leftarrow}(\nu))$, for all $\nu \in \mathbb{L}^Y$.

SL-NS denotes the category of all stratified \mathbb{L} -valued neighborhood spaces as objects and all continuous maps as morphisms.

Definition 8. [17, 22] Let $\Delta \subseteq \mathbb{L}^X$ such that the following are fulfilled:

- (LT1) $\top_X, \perp_X \in \Delta$;
- (LT2) $\nu_1, \nu_2 \in \Delta \Rightarrow \nu_1 \otimes \nu_2 \in \Delta$;
- (LT3) $\{\nu_j\}_{j \in J} \subseteq \Delta \Rightarrow \bigvee_{j \in J} \nu_j \in \Delta$;
- (SLT) $\nu \in \Delta, \alpha \in \mathbb{L} \Rightarrow \alpha_X * \nu \in \Delta$.

We call Δ an \mathbb{L} -valued topology on X if it satisfies (LT1)-(LT3), and the pair (X, Δ) is called an \mathbb{L} -valued topological space. If Δ satisfies (LT1)-(SLT) then we call it a stratified \mathbb{L} -valued topology on X and the pair (X, Δ) or X in short, if there is no confusion, is called a stratified \mathbb{L} -valued topological space; members of Δ are called open \mathbb{L} -valued sets or \mathbb{L} -valued subsets; the members of $\Theta(X) = \{\xi \in \mathbb{L}^X : \xi^c \text{ is open}\}$ are called closed \mathbb{L} -valued sets or \mathbb{L} -valued subsets, where ξ^c is the so-called quasi-complementation of ξ . Note that $\Theta(X)$ is closed under formation of arbitrary infs and finite sups. Furthermore, recall that the closure of $\nu \in \mathbb{L}^X$, denoted by $\bar{\nu}^X$ is defined as: $\bar{\nu}^X = \bigwedge \{\theta \in \Theta(X) : \nu \leq \theta\}$.

If (X, Δ) and (Y, Γ) are stratified \mathbb{L} -valued topological spaces, then a function $f: (X, \Delta) \rightarrow (Y, \Gamma)$ is said to be continuous if and only if for any $\sigma \in \Gamma$, $f^{\leftarrow}(\sigma) \in \Delta$. The category **SL-TOP** consists of all stratified \mathbb{L} -valued topological spaces as objects and all continuous maps between them as morphisms, while the category **L-TOP** consisting of all \mathbb{L} -valued topological spaces as objects and all continuous maps between them as morphisms.

Every stratified \mathbb{L} -valued topology Δ on X induces a stratified \mathbb{L} -valued neighborhood system $\mathfrak{N}_\Delta = (\mathfrak{N}_\Delta^x)$ as follows:

$$\mathfrak{N}_\Delta^x(\mu) = \bigvee \{\nu(x) : \nu \in \Delta, \nu \leq \mu\}, \text{ for all } \mu \in \mathbb{L}^X \text{ and } x \in X.$$

Conversely, every stratified \mathbb{L} -valued neighborhood system $\mathfrak{N} = (\mathfrak{N}^x)_{x \in X}$ on X induces a stratified \mathbb{L} -valued topology $\Delta_\mathfrak{N}$ on X :

$$\Delta_\mathfrak{N} = \{\nu \in \mathbb{L}^X : \nu(x) \leq \mathfrak{N}^x(\nu), \forall x \in X\}.$$

It follows that the interrelationship between \mathbb{L} -valued neighborhood system and \mathbb{L} -valued topologies can be viewed as:

$$\nu \in \Delta \Leftrightarrow \nu(x) \leq \mathfrak{N}^x(\nu), \forall x \in X \quad (\dagger).$$

As a consequence of (\dagger) it follows that the continuity between the objects in **SL-TOP**, and the continuity between objects in **SL-NS** are equivalent concept, cf. [18].

3. \mathbb{L} -valued topological groups and Kent convergence groups

We consider $\mathbb{L} = (\mathbb{L}, \leq, *, \otimes = *)$ an enriched cl -premonoid, where $*$ is a \mathbb{GL} -monoid operation. Let the category of groups and group homomorphisms be denoted by **GRP**.

Definition 9. Let $(X, \cdot) \in |\mathbf{GRP}|$ and $(X, \Delta) \in |\mathbf{S\mathbb{L}-TOP}|$. Then the triple (X, \cdot, Δ) is called a stratified \mathbb{L} -valued topological group if and only if the conditions below are fulfilled:

(LTGM) the mapping $m: (X \times X, \Delta \times \Delta) \rightarrow (X, \Delta), (x, y) \mapsto xy$ is continuous ;

(LTGI) the mapping $j: (X, \Delta) \rightarrow (X, \Delta), x \mapsto x^{-1}$ is continuous.

The category of all stratified \mathbb{L} -valued topological groups and continuous group homomorphisms is denoted by **S \mathbb{L} -TOPGRP**.

Definition 10. [3] Let $(X, \cdot) \in |\mathbf{GRP}|$ and $(X, \mathfrak{N} = (\mathfrak{N}^x)_{x \in X}) \in |\mathbf{S\mathbb{L}-NS}|$.

Then the triple $(X, \cdot, \mathfrak{N} = (\mathfrak{N}^x)_{x \in X})$ is called a stratified \mathbb{L} -valued neighborhood group if and only if

(LNGM) $\mathfrak{N}^{xy} \leq \mathfrak{N}^x \odot \mathfrak{N}^y$, and (LNGI) $\mathfrak{N}^{x^{-1}} \leq (\mathfrak{N}^x)^{-1}$ are satisfied, where for any $\xi \in \mathbb{L}^G$: $\mathfrak{N}^x \odot \mathfrak{N}^y(\xi) = m^{\Rightarrow}(\mathfrak{N}^x \times \mathfrak{N}^y)(\xi) = \bigvee \{ \mathfrak{N}^x(\xi_1) \wedge \mathfrak{N}^y(\xi_2) : \xi_1, \xi_2 \in \mathbb{L}^X, \xi_1 \times \xi_2 \leq m^{\leftarrow}(\xi) \}$.

A stratified \mathbb{L} -valued neighborhood system on a group X is said to be compatible with the group structure of X if and only if the group operations are continuous; i.e., conditions (LNGM) and (LNTGI) are fulfilled.

The category **S \mathbb{L} -NS** consists of all stratified \mathbb{L} -valued neighborhood groups as objects and continuous group homomorphisms as morphisms.

Example 1. Let $(G, \cdot) \in |\mathbf{GRP}|$, and $\mathfrak{N}^i: \mathbb{L}^X \rightarrow \mathbb{L}$ defined by $\mathfrak{N}^i = \bigwedge_{x \in G} [x]$. Then the triple $(G, \cdot, \mathfrak{N}^i)$ is a stratified \mathbb{L} -valued neighborhood group, called indiscrete stratified \mathbb{L} -valued neighborhood group.

Example 2. Let $(G, \cdot) \in |\mathbf{GRP}|$, and $\mathfrak{N}^d: \mathbb{L}^X \rightarrow \mathbb{L}$ defined by $\mathfrak{N}^{xd}(\nu) = \nu(x)$. Then the triple $(G, \cdot, \mathfrak{N}^d)$ is a stratified \mathbb{L} -valued neighborhood group, called discrete stratified \mathbb{L} -valued neighborhood group.

Lemma 2. [3] Let $(G, \cdot, \Delta) \in |\mathbf{S\mathbb{L}-TOPGRP}|$, and $a \in G$. Then the translations (left and right) $\mathcal{L}_a: (G, \cdot, \Delta) \rightarrow (G, \cdot, \Delta), g \mapsto ag$, and $\mathcal{L}_x: (G, \cdot, \Delta) \rightarrow (G, \cdot, \Delta), g \mapsto ga$ are homeomorphisms. Also the mapping $\mathcal{C}_a: (G, \cdot, \Delta) \rightarrow (G, \cdot, \Delta), g \mapsto gag^{-1}$ the inner automorphism is an isomorphism.

Definition 11. [20, 27] A Kent convergence structure q on X is a subset $q \subseteq \mathbb{F}(X) \times X$ such that the following conditions are satisfied:

(C1) $x \in q(\dot{x}), \forall x \in X$, where \dot{x} denotes the ordinary principal filter on X generated by the singleton $\{x\}$;

(C2) $\mathbb{F}, \mathbb{G} \in \mathbb{F}(X), \mathbb{F} \subseteq \mathbb{G}, x \in q(\mathbb{F})$ implies $x \in q(\mathbb{G})$;

(C3) $x \in q(\mathbb{F})$ implies $x \in q(\mathbb{F} \cap \dot{x})$.

Note that in [4], [6] and [7] the above notion is called a local filter convergence structure q on X , however.

A mapping $f: (X, q) \rightarrow (X', q')$ is called continuous if for all $\mathbb{F} \in \mathbb{F}(X)$ and $x \in X, x \in$

$q(\mathbb{F})$ implies $f(x) \in q(f(\mathbb{F}))$. The category of all Kent convergence spaces and continuous mapping is denoted by **KCONV**. The category **KCONV** is a strong topological universe, cf. [10, 28].

The pair (X, q) is called a limit space if conditions (C1), (C2) and (C4): $\forall \mathbb{F}, \mathbb{G} \in \mathbb{F}(X)$, $x \in q(\mathbb{F})$ and $x \in q(\mathbb{G})$ implies $x \in q(\mathbb{F} \cap \mathbb{G})$.

The category of limit spaces is denoted by **LIM**. A limit structure q on X is called a principal limit structure on X if and only if for every $x \in X$ there exists a unique filter $\mathbb{U}_x \in \mathbb{F}(X)$ such that the following relation holds:

$$q = \{(\mathbb{F}, x) \in \mathbb{F}(X) \times X : \mathbb{U}_x \subseteq \mathbb{F}\}.$$

The category of all principal limit spaces and continuous mappings is denoted by **pLIM**.

Remark 2. It is important to mention here that the categories of closure spaces, **CLS**, and **LIM** with principal limit structures are isomorphic, cf. [28], we are not interested at this stage to carry out research in this direction, and postpone it for further investigation.

Definition 12. [27] Let $(G, \cdot) \in |\mathbf{GRP}|$ and $(G, q) \in |\mathbf{KCONV}|$ (resp. $(G, q) \in |\mathbf{LIM}|$). Then the triple $(G, \cdot, q) \in |\mathbf{KCONVGRP}|$ (resp. $(G, \cdot, q) \in |\mathbf{LIMGRP}|$) if the following are fulfilled:

(CGM) $x \in q(\mathbb{F})$ and $y \in q(\mathbb{G})$ implies $xy \in q(\mathbb{F} \odot \mathbb{G})$;

(CGI) $x \in q(\mathbb{F})$ implies $x^{-1} \in q(\mathbb{F}^{-1})$.

The category of all Kent convergence groups and group homomorphisms is denoted by **KCONVGRP** (resp. the category of all limit groups and group homomorphisms is denoted by **LIMGRP**).

Given a stratified L -topological space $(X, \Delta_{\mathfrak{N}})$ with the corresponding \mathbb{L} -neighborhood system \mathfrak{N} . Then a filter \mathbb{F} is said to be convergent to a point $x \in X$ (we denoted it as $x \in q_{\Delta_{\mathfrak{N}}}(\mathbb{F})$) with respect to $\Delta_{\mathfrak{N}}$ if and only if for all $\nu \in \mathbb{L}^X$ the following holds:

$$\mathfrak{N}_x(\nu) \leq \bigvee_{F \in \mathbb{F}} \left(\bigwedge_{y \in F} \nu(y) \right).$$

Lemma 3. Let $(G, \cdot, \Delta_{\mathfrak{N}}) \in |\mathbf{SL-TOPGRP}|$, where Δ is a stratified L -valued topology on G and \mathfrak{N} is a corresponding \mathbb{L} -valued neighborhood system. Then $(G, \cdot, q_{\Delta_{\mathfrak{N}}}) \in |\mathbf{KCONVGRP}|$.

Proof. Let $(G, \cdot, \Delta_{\mathfrak{N}}) \in |\mathbf{SL-TOPGRP}|$. Then in view of the Lemma 5.4.1[18], we only need to Check the conditions (CGM) and (CGI).

(CGM) Let for $\mathbb{F}, \mathbb{G} \in \mathbb{F}(G)$ and $x, y \in G$, $x \in q_{\Delta_{\mathfrak{N}}}(\mathbb{F})$ and $y \in q_{\Delta_{\mathfrak{N}}}(\mathbb{G})$. Then for any $\nu, \mu \in \mathbb{L}^G$: $\mathfrak{N}_x(\nu) \leq \bigvee_{F \in \mathbb{F}} \bigwedge_{y_1 \in F} \nu(y_1)$, and $\mathfrak{N}_y(\mu) \leq \bigvee_{G \in \mathbb{G}} \bigwedge_{y_2 \in G} \mu(y_2)$. Thus, for any $\sigma \in \mathbb{L}^G$,

$$\begin{aligned} \mathfrak{N}_{xy}(\sigma) &\leq \bigvee \{ \mathfrak{N}_x(\nu) * \mathfrak{N}_y(\mu) : \nu(x) * \mu(y) \leq \sigma(xy) \} \leq \\ &\bigvee_{\nu(x) * \mu(y) \leq \sigma(xy)} \bigvee_{F, G \in \mathbb{F} \odot \mathbb{G}} \bigwedge_{y_1 \in F, y_2 \in G} \nu(x) * \mu(y) \leq \bigvee_{F, G \in \mathbb{F} \odot \mathbb{G}} \bigwedge_{xy \in F \cdot G} \sigma(xy) \end{aligned}$$

This implies that $\mathfrak{N}_{xy}(\sigma) \leq \bigvee_{F,G \in \mathbb{F} \odot \mathbb{G}} \bigwedge_{z \in F.G} \sigma(xy)$, i.e., $xy \in q_{\Delta_{\mathfrak{N}}}(\mathbb{F} \odot \mathbb{G})$.

(CGI) Let $\mathbb{F} \in \mathbb{F}(G)$, and $x \in X$. Then by invoking (†) in conjunction with the Lemma 5.4.1[18], if we consider $x \in q_{\Delta_{\mathfrak{N}}}(\mathbb{F})$, then for any $\nu \in \mathbb{L}^G$, we have $\mathfrak{N}_x(\nu) \leq \bigvee_{F \in \mathbb{F}} \left(\bigwedge_{y \in F} \nu(y) \right)$.

Now due to the continuity of j , we have

$$\mathfrak{N}_{x^{-1}}(\nu) \leq \mathfrak{N}_x(\nu^{-1}) \leq \bigvee_{F \in \mathbb{F}} \left(\bigwedge_{y \in F} \nu^{-1}(y) \right) = \bigvee_{F^{-1} \in \mathbb{F}^{-1}} \left(\bigwedge_{y^{-1} \in F^{-1}} \nu(y^{-1}) \right).$$

That is, $\mathfrak{N}_{x^{-1}}(\nu) \leq \bigvee_{F^{-1} \in \mathbb{F}^{-1}} \left(\bigwedge_{y^{-1} \in F^{-1}} \nu(y^{-1}) \right)$ implying $x^{-1} \in q_{\Delta_{\mathfrak{N}}}(\mathbb{F}^{-1})$.

Remark 3. Referring to the pp. 175 [18], one can observe that given a Kent convergence structure q on X , then q induces a stratified \mathbb{L} -valued topology $\widehat{\Delta}_q$ in the following way:

$$\widehat{\Delta}_q = \{ \sigma \in \mathbb{L}^X : \sigma(x) \leq \bigvee_{A \in \mathbb{F}} \left(\bigwedge_{z \in A} \sigma(z) \right), \forall \mathbb{F} \in \mathbb{F}(X), x \in q(\mathbb{F}) \}$$

From Lemma 5.4.2[18], it follows that there is a functor $\mathfrak{G}: \mathbf{KCONV} \rightarrow \mathbf{S\mathbb{L}\text{-TOP}}$, where $\mathfrak{G}(X, q) = (X, \widehat{\Delta}_q)$ and $\mathfrak{G}(f) = f$.

Lemma 4. Let $(G, \cdot, q) \in |\mathbf{KCONVGRP}|$. Then $(G, \cdot, \widehat{\Delta}_q) \in |\mathbf{S\mathbb{L}\text{-TOPGRP}|}$.

Proof. Let $(G, \cdot, q) \in |\mathbf{KCONVGRP}|$. Note that the product \mathbb{L} -valued topology on $\widehat{\Delta}_q \times \widehat{\Delta}_q$ is the initial \mathbb{L} -valued topology with respect to the projects $pr_1: X \times X \rightarrow X, (x, y) \mapsto x$, and $pr_2: X \times X \rightarrow X, (x, y) \mapsto y$. Further note that $\widehat{\Delta}_q \times \widehat{\Delta}_q = \{ (\nu^1 \cdot pr_1) * (\nu^2 \cdot pr_2) : \nu^1, \nu^2 \in \widehat{\Delta}_q \}$ is a base for the product \mathbb{L} -topology on $X \times X$, where the \mathbb{L} -set can be given by: $\mu_0 := \bigvee_{i \in I} (\nu_i^1 \cdot pr_1) * (\mu_i^2 \cdot pr_2)$, and $\nu_i^1, \mu_i^2 \in \widehat{\Delta}_q$. Thus, we have for any $\nu \in \widehat{\Delta}_q$ and $(x, y) \in X \times X$, and due to the property of $*$ in \mathbb{L} :

$$\begin{aligned} \nu(xy) &= m^{\leftarrow}(\nu)(x, y) = \bigvee_{i \in I} [(pr_1^{\leftarrow}(\nu_i^1)(x, y)) * (pr_2^{\leftarrow}(\mu_i^2)(x, y))] \quad (\nu_i^1, \mu_i^2 \in \widehat{\Delta}_q). \\ &= \bigvee_{i \in I} [\nu_i^1(x) * \mu_i^2(y)], \quad (\nu_i^1, \mu_i^2 \in \widehat{\Delta}_q). \\ &\leq \bigvee_{i \in I} \left[\bigvee_{A \in \mathbb{F}} \left(\bigwedge_{z_1 \in A} \nu_i^1(z_1) \right) * \bigvee_{B \in \mathbb{G}} \left(\bigwedge_{z_2 \in B} \mu_i^2(z_2) \right) \right] \\ &\leq \bigvee_{i \in I} \left[\bigvee_{A.B \in \mathbb{F} \odot \mathbb{G}} \bigwedge_{z_1 z_2 \in A.B} \left(\nu_i^1(z_1) * \mu_i^2(z_2) \right) \right] \\ &= \left[\bigvee_{A.B \in \mathbb{F} \odot \mathbb{G}} \bigwedge_{z_1 z_2 \in A.B} \nu(z_1 z_2) \right] \end{aligned}$$

That is, $\nu(xy) \leq \left[\bigvee_{H \in \mathbb{F} \odot \mathbb{G}} \bigwedge_{z_1 z_2 \in H} \nu(z_1 z_2) \right]$ and $xy \in q(\mathbb{F} \odot \mathbb{G})$ due to the condition (CGM) implying $m^{\leftarrow}(\nu) \in \widehat{\Delta}_q \times \widehat{\Delta}_q$. This proves condition (LTGM).

Now let $x \in q(\mathbb{F})$ for any $\mathbb{F} \in \mathbb{F}(G)$ and let $\nu \in \widehat{\Delta}_q$. Then we have

$$j^{\leftarrow}(\nu)(x) = \nu(j(x)) \leq \bigvee_{A \in \mathbb{F}} \left(\bigwedge_{z_2 \in j(A)} \nu(z_2) \right) = \bigvee_{A^{-1} \in \mathbb{F}^{-1}} \left(\bigwedge_{z_1 \in A^{-1}} j^{\leftarrow}(\nu)(z_1) \right),$$

that is, $j^{\leftarrow}(\nu)(x) \leq \bigvee_{A^{-1} \in \mathbb{F}^{-1}} \left(\bigwedge_{z_1 \in A^{-1}} j^{\leftarrow}(\nu)(z_1) \right)$; and $x^{-1} \in q(\mathbb{F}^{-1})$ because of the condition (CGI). These together imply that $j^{\leftarrow}(\nu) \in \widehat{\Delta}_q$, this proves (LTGI).

Theorem 1. The functor $\mathfrak{F}: \mathbf{S\mathbb{L}\text{-TOPGRP}} \rightarrow \mathbf{KCONVGRP}$ as defined below

$$\mathfrak{F} : \begin{cases} \mathbf{S\mathbb{L}\text{-TOPGRP}} & \longrightarrow & \mathbf{KCONVGRP} \\ (G, \cdot, \Delta_{\mathfrak{N}}) & \longmapsto & (G, \cdot, q_{\Delta_{\mathfrak{N}}}) \\ f & \longmapsto & f \end{cases}$$

has a left adjoint.

Proof. In view of Lemma 3 in conjunction with Lemma 5.4.1 [18], $\mathfrak{F}: \mathbf{SL-TOPGRP} \longrightarrow \mathbf{KCONVGRP}$ is a functor. Define $\mathfrak{G}: \mathbf{KCONVGRP} \longrightarrow \mathbf{SL-TOPGRP}$ by

$$\mathfrak{G} : \begin{cases} \mathbf{KCONVGRP} & \longrightarrow & \mathbf{SL-TOPGRP} \\ (G, \cdot, q) & \longmapsto & (G, \cdot, \widehat{\Delta}_q) \\ f & \longmapsto & f \end{cases}$$

Then from Lemma 4 in conjunction with Lemma 5.4.2 [18] that \mathfrak{G} is a functor since in both the cases the group homomorphism structures remain unchanged. That the functor \mathfrak{G} is a left adjoint since in both the cases group homomorphism structures remain unchanged. That the functor \mathfrak{G} is a left adjoint to \mathfrak{F} is an immediate consequence of the Proposition 5.4.3 [18].

4. Enriched lattice-valued subgroup of a group and enriched lattice-valued neighborhood groups

Definition 13. Let $\mathbb{L} = (\mathbb{L}, \leq, \wedge, *)$ be an enriched cl-premonoid, $(G, \cdot) \in |\mathbf{GRP}|$. Then an \mathbb{L} -set $\mu: G \longrightarrow \mathbb{L}$ is called an \mathbb{L} -valued subgroup of a group G if and only if the following conditions are fulfilled:

- (LG1) $\mu(e) = \top$;
- (LG2) $\mu(g) * \mu(h) \leq \mu(gh), \forall g, h \in G$;
- (LG3) $\mu(g) \leq \mu(g^{-1})$.

Then the pair (G, \cdot, μ) is called an \mathbb{L} -valued subgroup space. Let (H, \cdot, ξ) be another \mathbb{L} -valued subgroup of a group H . Define a mapping between \mathbb{L} -valued subgroup spaces, $f: (G, \cdot, \mu) \longrightarrow (H, \cdot, \xi)$ such that

$$\mu(g) \leq \xi(f(g)), \forall g \in G \quad (\ddagger)$$

The category of all \mathbb{L} -valued subgroup spaces and all group homomorphisms satisfying (\ddagger) is denoted by $\mathbb{L}\text{-GRP}$. Sometime we denote the set of \mathbb{L} -valued subgroups of a group G by $\mathbb{L}(G)$.

Example 3. [3] Let $\mathbb{L} = ([0, 1], \leq, \wedge, *)$ be an enriched cl-premonoid, where $*$ is a t-norm on $[0, 1]$. Let G be the cyclic group C_n of order n ($n \geq 1$) with a as the generator; specifically, $C_n = \{e, a, a^2, \dots, a^{n-1}; a^n = e\}$ with respect to multiplication \cdot .

Define $\mu: G \rightarrow [0, 1]$ by

$$\mu(x) = \begin{cases} 1, & \text{if } x = e; \\ \frac{1}{n}, & \text{otherwise.} \end{cases}$$

Then (G, \cdot, μ) is an enriched lattice-valued subgroup space. In fact, for (LG1) $\mu(e) = 1$ while (LG3) follows from the definition. For (LG2), consider $x, y \in G$ with $x \neq e$ and $y \neq e$, then $\mu(x) * \mu(y) = \frac{1}{n} * \frac{1}{n} \leq \frac{1}{n} * 1 = \frac{1}{n}$ implying $\mu(x) * \mu(y) \leq \mu(xy)$; other choices follow similarly. Hence μ is an enriched lattice-valued subgroup of the group G .

Remark 4. In [33], C. L. Walker pointed out that for a category of fuzzy subsets $\mathbf{F} = \mathbf{Set}(I)$ where all objects are (X, ν) , $X \in |\mathbf{Set}|$, with $\nu: X \rightarrow I$ - a mapping from X to the unit interval. The morphisms \mathbf{F} are all mappings $f: (X, \nu) \rightarrow (Y, \mu)$ satisfying $\nu(x) \leq \mu(f(x))$. Furthermore, note that in [14], J. Goguen, defined the category $\mathbf{SET}(\mathbb{L})$ having objects the pair (X, ν) , where $\nu: X \rightarrow \mathbb{L}$, and morphisms $f: (X, \nu) \rightarrow (Y, \mu)$ such that $\nu(x) \leq \mu(f(x))$ holds. L. Stout [32] argued that this category $\mathbf{SET}(\mathbb{L})$ has initial structure and is cartesian closed. The initial structure is given as: for a family of mappings $(f_j: X \rightarrow (Y_j, \mu_j))_{j \in J}$, $\nu(x) = \bigwedge_j \mu_j(f_j(x))$ gives the initial structure on X . The cartesian closed structure is obtained as: $(\mathcal{C}(X, Y), \nabla)$, where $\nabla(f) = \bigwedge_{x \in X} [\nu(x) \rightarrow \mu(f(x))]$, where for all $(f: (X, \nu) \rightarrow (Y, \mu)) \in \mathcal{C}(X, Y)$, and the implication \rightarrow is given by: $\nu(x) \rightarrow \mu(f(x)) = \bigvee \{ \lambda: \lambda \wedge \nu(x) \leq \mu(f(x)) \}$.

Lemma 5. \mathbb{L} -GRP has initial structure where the underlying forgetful functor is given by $\mathfrak{F}: \mathbb{L}\text{-GRP} \rightarrow \mathbf{GRP}$.

Proof. Consider a group (G, \cdot) and a family of mappings $(f_j: G \rightarrow (H_j, \mu_j))_{j \in J}$, where each $f_j: G \rightarrow H_j$ is a group homomorphism, μ_j is a subgroup of H_j , for each $j \in J$. Then the structure on μ on G is given by $\nu(g) = \bigwedge_j \mu_j(f_j(g)) (= \bigwedge_j f_j^{\leftarrow}(\mu_j)(g))$, for all $g \in G$, note that for each $j \in J$, $f_j^{\leftarrow}(\mu_j)$ is also an \mathbb{L} -subgroup of G , and the arbitrary intersection ν is also an \mathbb{L} -subgroup of G , and hence $(G, \cdot, \nu) \in |\mathbb{L}\text{-GRP}|$. Let $(Z, \cdot, \varrho) \in |\mathbb{L}\text{-GRP}|$, we prove that the mapping $\varphi: (Z, \cdot, \varrho) \rightarrow (G, \cdot, \nu)$ a group homomorphism is an $\mathbb{L}\text{-GRP}$ -morphism if and only if $f_j \circ \varphi: (Z, \cdot, \varrho) \rightarrow (H_j, \cdot, \mu_j)$ is an $\mathbb{L}\text{-GRP}$ -morphism. We only show $g: (Z, \cdot, \varrho) \rightarrow (G, \cdot, \nu)$ is an $\mathbb{L}\text{-GRP}$ -morphism. So, for any $z \in Z$, $\varrho(z) \leq \mu_j(f_j(\varphi(z))) = \bigwedge_{j \in J} f_j^{\leftarrow}(\mu_j)(\varphi(z)) = \nu(\varphi(z))$, i.e., $\varrho(z) \leq \nu(\varphi(z))$.

Theorem 2. Let $\mathbb{L} = (\mathbb{L}, \leq, * = \wedge)$ be a complete Heyting algebra, and (G, \cdot, ν) be an \mathbb{L} -valued subgroup space and $\mathcal{T}(G) = \{f: (G, \nu) \rightarrow (G, \nu); f \text{ is bijective and both } f \text{ and } f^{-1} \text{ satisfy } (\ddagger)\}$. Then $(\mathcal{T}(G), \cdot, \nabla)$ is an \mathbb{L} -subgroup space, where $(fg)(x) = f(x)g(x)$ and $f^{-1}(x) = (f(x))^{-1}$.

Proof. Clearly $(\mathcal{T}(G), \cdot)$ is a group under composition. Define $\nabla(f) = \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(f(x))]$, $\forall f \in \mathcal{T}(G)$ (a)

and $\nabla^{(-1)}(f) = \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(f^{-1}(x))]$, $\forall f \in \mathcal{T}(G)$ (b)

Combining (a) and (b) it follows upon using Proposition 1(7) that $\nabla(f) = \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(f(x)) \wedge \nu(f^{-1}(x))]$. Then clearly (LG1) and (LG3) are true upon using Proposition 1(7) and (LG2), i.e., $\nabla(id_G) = \top$, and $\nabla(f) \leq \nabla(f^{-1})$; we only look at (LG2).

For, let $f, g \in \mathcal{T}(X)$, then we have $\nabla(f) \wedge \nabla(g) = \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(f(x)) \wedge \nu(f^{-1}(x))] \wedge \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(g(x)) \wedge \nu(g^{-1}(x))] \leq \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(f(x)) \wedge \nu(g(x)) \wedge \nu(g^{-1}(x)) \wedge \nu(f^{-1}(x))] \leq \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(f(x)g(x)) \wedge \nu(g^{-1}(x)f^{-1}(x))] = \bigwedge_{x \in G} [\nu(x) \rightarrow \nu(fg(x)) \wedge \nu((fg)^{-1}(x))] = \nabla(fg)$.

Definition 14. [23, 25] An \mathbb{L} -valued subgroup is called \mathbb{L} -valued normal subgroup if for all $x, y \in G$ if it satisfies one of the following equivalent conditions:

- (1) $\nu(xy) = \nu(yx)$;
- (2) $\nu(xy x^{-1}) \geq \nu(y)$;
- (3) $\nu(xy x^{-1}) = \nu(y)$.

Definition 15. A mapping $\ell: \mathbb{L}^X \rightarrow \mathbb{L}^X$ is said to be an \mathbb{L} -valued closure operation on X if the following conditions hold for every $\nu, \mu \in \mathbb{L}^X$:

- (1) $\nu \leq \ell(\nu)$;
- (2) $\nu \leq \mu$ implies $\ell(\nu) \leq \ell(\mu)$;
- (3) $\ell(\ell(\nu)) = \ell(\nu)$;
- (4) $\ell(\top_\emptyset) = \perp$.

The pair (X, ℓ) is called an \mathbb{L} -valued closure space and $\nu \in \mathbb{L}^X$ is called closed if $\nu = \ell(\nu)$. Note that (2) implies $\ell(\nu) \vee \ell(\mu) \leq \ell(\nu \vee \mu)$, for any $\nu, \mu \in \mathbb{L}^X$.

The category of all \mathbb{L} -valued closure spaces and all closure preserving mappings, i.e., mappings $f: (X, \ell) \rightarrow (Y, \ell)$ that satisfy $f^\rightarrow(\ell(\nu)) \leq \ell(f^\rightarrow(\nu))$ for all $\nu \in \mathbb{L}^X$, is denoted by $\mathbb{L}\text{-CLS}$.

Lemma 6. We have the following forgetful functor forgetting \mathbb{L} -valued closure structure:

$$\mathfrak{U} : \begin{cases} \mathbb{L}\text{-CLS} & \longrightarrow & \mathbf{SET}(\mathbb{L}) \\ (X, \ell) & \longmapsto & (X, \nu) \\ f & \longmapsto & f \end{cases}$$

where $\mathfrak{U}((X, \ell)) = (X, \nu)$ and for $f: X \rightarrow Y$, $\mathfrak{U}(f) = f$, $f^\rightarrow: \mathbb{L}^X \rightarrow \mathbb{L}^Y$, and $\mathfrak{U}(f)$ yields an $\mathbf{SET}(\mathbb{L})$ -morphism.

Let $X \in |\mathbf{SET}|$ and let $\Omega \subset \mathbb{L}^X$ be a collection of \mathbb{L} -subsets of X . Then we call Ω a lattice-valued Moore collection if every intersection of members of Ω belongs to Ω , i.e., given a family $(\nu_j)_{j \in J}$ of \mathbb{L} -subsets: $\forall j \in J, \nu_j \in \Omega \implies \bigwedge_{j \in J} \nu_j \in \Omega$. If Ω is a lattice-valued Moore collection containing \top_\emptyset , then if $\ell(\mu)_\Omega = \bigwedge\{\nu \in \Omega: \mu \leq \nu, \nu \text{ is } \mathbb{L}\text{-valued closed set}\}$, i.e. if $\ell(\mu)$ is the intersection of all \mathbb{L} -valued closed sets that contain μ , then ℓ is an \mathbb{L} -valued closure operator. We refer to Birkhoff [9], and Schechter [31], for the classical notion of Moore collection.

Example 4. \mathbb{L} -valued subgroups of a group (G, \cdot) form a lattice-valued Moore collection; this is so, since arbitrary intersection of \mathbb{L} -valued subgroups is again an \mathbb{L} -valued subgroup, cf. [11], pp. 115. In fact, if we let $\mu = \bigwedge_{j \in J} \nu_j$, then we can easily verify the Definition 13. In fact, (LG1) $\mu(e) = \bigwedge \nu_j(e) = \top$ for all $j \in J$; (LG2) upon using Proposition 1(10), we have: $\mu(x) * \mu(y) = (\bigwedge_{j \in J} \nu_j(x)) * (\bigwedge_{j \in J} \nu_j(y)) \leq \bigwedge_{j \in J} (\nu_j(x) * \nu_j(y)) \leq \bigwedge_{j \in J} \nu_j(xy) = \mu(xy)$, so, $\mu(x) * \mu(y) \leq \mu(xy)$; (LG3) $\mu = \bigwedge_{j \in J} (\nu_j(x)) \leq \bigwedge_{j \in J} (\nu_j(x^{-1})) = \mu(x^{-1})$. Also, if $\mu \in \mathbb{L}^G$, then \mathbb{L} -valued closure of μ is the subgroup generated by μ . This can be given as:

$$\langle \ell(\mu) \rangle = \bigwedge\{\nu: \mu \leq \nu, \nu \text{ is closed } \mathbb{L}^G\text{-valued subgroup of } G\},$$

the \mathbb{L} -valued subgroup that contains μ .

In view of the Theorem 5.2.6[11], normal \mathbb{L} -valued subgroup of the group G form a lattice-valued Moore collection, and in particular, $\ell(\mu)$, $\mu \in \mathbb{L}^G$ is the normal \mathbb{L} -valued subgroup generated by μ . More precisely, $\langle \ell(\mu) \rangle = \bigwedge \{ \nu : \mu \leq \nu, \nu \text{ is closed normal } \mathbb{L}^G\text{-valued subgroup of } G \}$,

Theorem 3. $\mathbb{L}\text{-CLS}$ is a topological category.

Proof. Note that the objects of $\mathbb{L}\text{-CLS}$ are structured sets and the composition of closure preserving mappings is closure preserving.

Consider X is a set, $(Y_j, \ell^j)_{j \in J}$ a family of \mathbb{L} -valued closure spaces and a source $\mathcal{S} = (f_j : X \rightarrow (Y_j, \ell^j))_{j \in J}$ of family of functions, then

$$\Omega = \{ \omega \in \mathbb{L}^X : \omega = \bigwedge_{j \in J} f_j^{\leftarrow}(\omega_j), \forall \omega_j = \ell_j(\omega_j), j \in J \}$$

is a lattice-valued Moore family which contains \top_\emptyset . Then Ω induces an \mathbb{L} -valued closure operation on X given by: $\ell(\mu)_\Omega = \bigwedge \{ \omega \in \Omega : \mu \leq \omega \}$, for all $\mu \in \mathbb{L}^X$. Now let $(Z, \ell) \in |\mathbb{L}\text{-CLS}|$, and $g : Z \rightarrow X$ be a function such that $f_j \circ g : (Z, \ell) \rightarrow (Y_j, \ell_j)$ is closure preserving mapping for all $j \in J$. If $\mu \in \mathbb{L}^X$ is a $^{-\Omega}$ closed, then $\mu \in \Omega$ and thus $\mu = \bigwedge_{j \in J} f_j^{\leftarrow}(\omega_j)$ where $\omega_j = \ell_j(\omega_j)$ in (Y_j, ℓ_j) . In view of Proposition 1.2(5) [22], we have:

$$g^{\leftarrow}(\mu) = g^{\leftarrow} \left(\bigwedge_j f_j^{\leftarrow}(\omega_j) \right) = \bigwedge_j g^{\leftarrow}(f_j^{\leftarrow}(\omega_j)) = \bigwedge_j (f_j \circ g)^{\leftarrow}(\omega_j)$$

This implies $(f_j \circ g)^{\leftarrow}(\omega_j)$ is closed in (Z, ℓ) implying $g^{\leftarrow}(\mu)$ is closed in (Z, ℓ) .

Remark 5. Every \mathbb{L} -valued topological space (X, Δ) is an \mathbb{L} -valued closure space with the closure operation defined by: $\ell(\nu) = \overline{\nu}^{(X, \Delta)} = \overline{\nu}^X$ for every $\nu \in \mathbb{L}^X$. Also, every mapping $f : (X, \Delta) \rightarrow (Y, \Gamma)$ continuous if and only if it is closure preserving with respect to the induced \mathbb{L} -valued closure operations. In fact, if $\nu \in \mathbb{L}^X$, then in view of the Proposition 1.4 [22], $f^{\rightarrow}(\ell(\nu)) = f^{\rightarrow}(\overline{\nu}^X) \leq \overline{f^{\rightarrow}(\nu)}^Y = \ell(f^{\rightarrow}(\nu))$, i.e., $f^{\rightarrow}(\ell(\nu)) \leq \ell(f^{\rightarrow}(\nu))$, meaning f is closure preserving. Conversely, let $\nu \in \mathbb{L}^X$ and f be closure preserving, then $f^{\rightarrow}(\overline{\nu}^X) = f^{\rightarrow}(\ell(\nu)) \leq \ell(f^{\rightarrow}(\nu)) = \overline{f^{\rightarrow}(\nu)}^Y$, i.e., $f^{\rightarrow}(\overline{\nu}^X) \leq \overline{f^{\rightarrow}(\nu)}^Y$ meaning the mapping $f : (X, \Delta) \rightarrow (Y, \Gamma)$ is continuous by the Proposition 1.4 [22]. Thus we have the following.

Corollary 1. $\mathbb{L}\text{-TOP}$, the category of \mathbb{L} -valued topological spaces and continuous mappings is a full subcategory of the category $\mathbb{L}\text{-CLS}$

Definition 16. A triple (G, \cdot, ℓ) is called an \mathbb{L} -closure group if $(G, \cdot) \in |\mathbf{GRP}|$ and $(G, \ell) \in |\mathbb{L}\text{-CLS}|$ such the following are fulfilled:

(clGM) $\ell(\nu)(x) * \ell(\nu)(y) \leq \ell(\nu \cdot \nu)(xy)$, $\forall \nu \in \mathbb{L}^G$ and $\forall x, y \in G$;

(clGI) $\ell(\nu)(x) \leq \ell(\nu^{-1})(x^{-1})$, $\forall \nu \in \mathbb{L}^G$ and $x \in G$.

The category of all \mathbb{L} -valued closure groups and closure-preserving group homomorphisms is denoted by $\mathbb{L}\text{-CLGRP}$.

Remark 6. If we consider each $\nu \in \mathbb{L}(G)$, i.e., each $\nu \in \mathbb{L}^G$ is an \mathbb{L} -valued subgroup of the group G , then we obtain a category $\mathbb{L}\text{-CLGRP}^*$ of all \mathbb{L} -valued closure of \mathbb{L} -valued subgroups of G , and closure-preserving mappings. Then $\mathbb{L}\text{-CLGRP}^*$ is a subcategory of $\mathbb{L}\text{-CLGRP}$.

Theorem 4. $\mathbb{L}\text{-CLGRP}$ is a topological category.

Proof. Consider (G, \cdot) a group, and a source $\mathcal{S} = (f_j: (G, \cdot) \rightarrow (G_j, \cdot, \ell_j))_{j \in J}$ of family of functions, where for each $j \in J$, $f_j: G \rightarrow G_j$ is a group homomorphism, then

$$\Omega = \{\omega \in \mathbb{L}^G : \omega = \bigwedge_{j \in J} f_j^{\leftarrow}(\omega_j), \forall \omega_j = \ell_j(\omega_j), j \in J\}$$

In view of Theorem 3, we have (G, \cdot, ℓ) is an \mathbb{L} -valued closure space. We only verify (clGM). So we have:

$$\begin{aligned} \ell(\omega)(x) * \ell(\omega)(y) &= \bigwedge_{j \in J} f_j^{\leftarrow}(\omega_j)(x) * \bigwedge_{j \in J} f_j^{\leftarrow}(\omega_j)(y) \leq \bigwedge_{j \in J} f_j^{\leftarrow}(\omega_j) \odot f_j^{\leftarrow}(\omega_j)(xy) \\ &= \bigwedge_{j \in J} f_j^{\leftarrow}(\omega_j \odot \omega_j)(xy) \leq \ell(\omega \cdot \omega)(xy). \end{aligned}$$

Definition 17. [8, 19, 32] An \mathbb{L} -tolerance space is a pair (X, τ) , where $\tau: X \times X \rightarrow \mathbb{L}$ such that

(T1) $\tau(x, x) = \top, \forall x \in X$ (reflexivity);

(T2) $\tau(x, y) = \tau(y, x)$ (symmetry).

If, in addition τ satisfies (T3) $\tau(x, y) * \tau(y, z) \leq \tau(x, z)$, for any $x, y, z \in X$, then we speak of transitive tolerance relation which is essentially gives an \mathbb{L} -equivalence relation.

A mapping between \mathbb{L} -valued tolerance spaces (resp. transitive \mathbb{L} -valued tolerance spaces): $f: (X, \tau) \rightarrow (Y, \tau')$ is called \mathbb{L} -valued tolerance preserving if $\tau(x, y) \leq \tau'(f(x), f(y))$.

The category of all \mathbb{L} -valued tolerance spaces and \mathbb{L} -tolerance preserving mappings is denoted by $\mathbb{L}\text{-TOL}$ while $\mathbb{L}\text{-TranTOL}$ denotes the category of transitive \mathbb{L} -tolerance spaces.

For an \mathbf{MV} -valued algebra \mathbb{L} , given $\mathbb{L}\text{-TranTOL}$ a category of transitive \mathbb{L} -valued tolerance spaces and \mathbb{L} -valued tolerance preserving mappings, one can obtain a functor $\mathcal{A}: \mathbb{L}\text{-TOL} \rightarrow \mathbb{L}\text{-SET}$ where $\mathcal{A}(X, \tau) = (X, \tau\mathbb{D})$, $\mathbb{D}: X \rightarrow X \times X$ and $\mathcal{A}(f) = f$, here $\mathcal{A}(f)$ sends f to an \mathbb{L} -tolerance preserving mapping to $f: (X, \tau\mathbb{D}) \rightarrow (Y, \tau'\mathbb{D})$, i.e., $\tau\mathbb{D}(x) = \tau(x, x) \leq \tau'(f(x), f(x)) = \tau'\mathbb{D}(f(x))$, i.e., $\tau\mathbb{D}(x) \leq \tau'\mathbb{D}(f(x))$. Conversely, given $\mathbb{L}\text{-SET}$, one obtains a functor $\mathcal{B}: \mathbb{L}\text{-SET} \rightarrow \mathbb{L}\text{-TranTOL}$ as defined by: $\mathcal{B}(X, \nu) = (X, \tau := \nu \wedge \nu)$ and $\mathcal{B}(f) = f, \tau(x, y) = \nu(x) \wedge \nu(y) \leq \nu(f(x)) \wedge \nu(f(y)) = \tau(f(x), f(y))$.

In view of [11], pp 148, for a group (G, \cdot) , we consider a mapping $\varrho_L: \mathbb{L}^G \rightarrow \mathbb{L}^{G \times G}$ defined by: $\varrho_L(\nu)(x, y) = \nu(x^{-1}y)$, and analogously, $\varrho_R(\nu)(x, y) = \nu(xy^{-1})$. Then we have the following.

Lemma 7. Let $(G, \cdot) \in |\mathbf{GRP}|$, and the category $\mathbb{L}\text{-TranTOL}$ consists of morphisms $f: (G, \tau) \rightarrow (H, \varrho')$ which are \mathbb{L} -valued tolerance preserving such that each morphism is a group homomorphism. Then

$$\mathfrak{A} : \begin{cases} \mathbb{L}\text{-GRP} & \longrightarrow & \mathbb{L}\text{-TranTOL} \\ (G, \nu) & \longmapsto & (G, \varrho_L(\nu)) \\ f & \longmapsto & f \end{cases}$$

Proof. Let $\nu \in \mathbb{L}(G)$, then we have $\rho_L(\nu)(x, x) = \nu(x^{-1}x) = \nu(e) = \top$ which is (T1); for (T2), we apply Theorem 5.1.1(5)[11](see also, Theorem 1.2.2[24]) to get $\rho_L(\nu)(x, y) = \nu(x^{-1}y) = \nu((x^{-1}y)^{-1}) = \nu(y^{-1}x) = \rho_L(\nu)(y, x)$. Now for any $x, y, z \in X$, $\rho_L(\nu)(x, y) * \rho_L(\nu)(y, z) = \nu(x^{-1}y) * \nu(y^{-1}z) \leq \nu(x^{-1}yy^{-1}z) = \nu(x^{-1}z) = \rho_L(\nu)(x, z)$, which is (T3). To check the morphism part, we have for any $x, y \in G$ and $\nu \in \mathbb{L}(G)$: $\tau(x, y) = \rho_L(\nu)(x, y) = \nu(x^{-1}y) \leq \nu'(f(x^{-1}y)) = \nu'((f(x))^{-1}f(y)) = \rho_L(\nu)(f(x), f(y)) = \tau'(f(x), f(y))$, i.e., $\tau(x, y) \leq \tau'(\nu')(f(x), f(y))$.

Lemma 8. *Let $(G, \cdot) \in |\mathbf{GRP}|$, and the category $\mathbb{L}\text{-TranTOL}$ consists of morphisms $f: (G, \rho_L(\nu)) \rightarrow (H, \rho_L(\nu'))$ which are \mathbb{L} -valued tolerance preserving such that each morphism is a group homomorphism. Then*

$$\mathfrak{B} : \begin{cases} \mathbb{L}\text{-TranTOL} & \longrightarrow & \mathbb{L}\text{-GRP} \\ (G, \rho_L(\nu)) & \longmapsto & (G, \nu) \\ f & \longmapsto & f \end{cases}$$

Proof. Let $\nu \in \mathbb{L}^G$, and $(G, \rho_L(\nu)) \in |\mathbb{L}\text{-TranTOL}|$, it suffices to show that $\nu \in \mathbb{L}(G)$. Thus, for any $x \in X$, $\nu(e) = \nu(x^{-1}x) = \rho_L(\nu)(x, x) = \top$ which is (LG1). For (LG2) is obviously true while for (LG3), we have for any $x, y \in G$: $\nu(x) * \nu(y) = \nu(xe) * \nu(ey) = \rho_L(\nu)(x, e) * \rho_L(\nu)(e, y) \leq \rho_L(\nu)(x, y) = \nu(x^{-1}y)$, i.e., $\nu(x) * \nu(y) \leq \nu(x^{-1}y)$, this happens when we combine (LG2) and (LG3), cf. Theorem 5.1.3[11]. This shows that $\nu \in \mathbb{L}(G)$. For the morphism part, let $x \in G$ and $\nu \in \mathbb{L}^G$. Then $\nu(x) = \nu(ex) = \rho_L(\nu)(e, x) \leq \rho_L(\nu')(f(e), f(x)) = \nu'((f(e))^{-1}f(x)) = \nu'(f(ex)) = \nu'(f(x))$, i.e., $\nu(x) \leq \nu'(f(x))$.

5. Enriched latticed-valued subgroups on lattice-valued neighborhood groups

Let $\mathbb{L} = (\mathbb{L}, \leq, *)$ be a complete **MV**-valued algebra with square roots. If $(X, \mathfrak{N} = (\mathfrak{N}_x)_{x \in X})$ is a stratified \mathbb{L} -neighborhood space, then in view of [12] (page 13), and [18] (page 226), one can see that \mathfrak{N} induces a closure operator $\bar{\cdot} : \mathbb{L}^X \rightarrow \mathbb{L}^X$ given for any $x \in X$ and $\nu \in \mathbb{L}^X$ by

$$\bar{\nu}(x) = ([\mathfrak{N}_x](\nu \rightarrow \perp)) \rightarrow \perp.$$

Theorem 5. [3, 12, 18] (a) *Let $(X, \mathfrak{N} = (\mathfrak{N}_x)_{x \in X}) \in |\mathbf{SL-NS}|$. Then*

$$\bar{\nu}(x) = \bigvee \{ \mathcal{F}(\nu) : \mathcal{F} \in \mathcal{F}_{\mathbb{L}}^s(X), \mathcal{F} \geq \mathfrak{N}_x \}, \forall \nu \in \mathbb{L}^X, \text{ and } \forall x \in X.$$

(b) *Let $(G, \cdot, \mathfrak{N} = (\mathfrak{N}_x)_{x \in G}) \in |\mathbf{SL-NGRP}|$ and $\nu \in \mathbb{L}^G$ be an \mathbb{L} -valued subgroup of a group G . Then the \mathbb{L} -valued closure $\bar{\nu}$ of ν in (a) is an \mathbb{L} -valued subgroup of G .*

(c) *Let $(G, \cdot, \mathfrak{N}) \rightarrow (H, \cdot, \mathfrak{M})$ be continuous group homomorphism. Then $\bar{\nu}(x) \leq \overline{f^{\rightarrow}(\nu)}(f(x))$ for all $\nu \in \mathbb{L}^G$ and $x \in G$. Moreover, if $\nu \in \mathbb{L}^G$ is an \mathbb{L} -valued subgroup of G , then $\overline{f^{\rightarrow}(\nu)}$ is an \mathbb{L} -valued subgroup of H , and if $\mu \in \mathbb{L}^G$ is a \mathbb{L} -valued subgroup of H , then $f^{\leftarrow}(\mu)$ is an \mathbb{L} -valued subgroup of G .*

(d) *If $\nu \in \mathbb{L}^G$ is an \mathbb{L} -valued normal subgroup of a group G , then $\bar{\nu}$ is also an \mathbb{L} -valued*

normal subgroup of G .

(e) If $(G, \cdot, \mathfrak{N}) \rightarrow (H, \cdot, \mathfrak{M})$ is a continuous group homomorphism and $\mu \in \mathbb{L}^H$ is an \mathbb{L} -valued subgroup of H , then $f^{\leftarrow}(\mu)$ is an \mathbb{L} -valued subgroup of G .

Proof. (b) follows from the Theorem 5.1[3].

(c) Let $\nu \in \mathbb{L}^G$, and $x \in G$. Then since $\nu \leq f^{\leftarrow}(f^{\rightarrow}(\nu))$ due to Definition 6 (LF2), $\mathcal{F}(\nu) \leq \mathcal{F}(f^{\leftarrow}(f^{\rightarrow}(\nu))) = f^{\rightarrow}(\mathcal{F})(f^{\rightarrow}(\nu))$, and since $\mathfrak{M}_{f(x)} \leq f^{\rightarrow}(\mathfrak{N}_x)$ due to continuity of f , we have

$$\begin{aligned} \bar{\nu}(x) &= \bigvee \{ \mathcal{F}(\nu) : \mathcal{F} \in \mathcal{F}_{\mathbb{L}}^s(X), \mathcal{F} \geq \mathfrak{N}_x \} \leq \bigvee \{ f^{\rightarrow}(\mathcal{F})(f^{\rightarrow}(\nu)) : f^{\rightarrow}(\mathcal{F}) \in \mathcal{F}_{\mathbb{L}}^s(Y), f^{\rightarrow}(\mathcal{F})(f^{\rightarrow}(\nu)) \geq f^{\rightarrow}(\mathfrak{N}_x)(f^{\rightarrow}(\nu)) \} \\ &\leq \bigvee \{ f^{\rightarrow}(\mathcal{F})(f^{\rightarrow}(\nu)) : f^{\rightarrow}(\mathcal{F}) \in \mathcal{F}_{\mathbb{L}}^s(H), f^{\rightarrow}(\mathcal{F})(f^{\rightarrow}(\nu)) \geq \mathfrak{M}_{f(x)}(f^{\rightarrow}(\nu)) \} \\ &= \bigvee \{ \mathcal{G}(f^{\rightarrow}(\nu)) : \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(Y), \mathcal{G} \geq \mathfrak{M}_{f(x)} \} = \overline{f^{\rightarrow}(\nu)}(f(x)), \text{ i.e., } \bar{\nu}(x) \leq \overline{f^{\rightarrow}(\nu)}(f(x)). \end{aligned}$$

(d) Let $\nu \in \mathbb{L}^G$ be an \mathbb{L} -valued normal subgroup of a group G , and consider the mapping $\mathcal{C}_a : G \rightarrow G$ defined by $\mathcal{C}_a(g) = a^{-1}ga$; need to that $\bar{\nu}$ is also an \mathbb{L} -normal subgroup of G . Note that ν is \mathbb{L} -normal subgroup of G if and only if $\nu(aga^{-1}) = \nu(g)$. Now since the mapping \mathcal{C}_a is continuous, we have $\bar{\nu}(g) \leq \overline{\mathcal{C}_a(\nu)}(\mathcal{C}_a(g)) = \bigvee_{x \in \mathcal{C}_a^{-1}(g)} \bar{\nu}(x) = \bigvee_{\mathcal{C}_a(x)=g} \bar{\nu}(x) = \bar{\nu}(aga^{-1})$, i.e., $\bar{\nu}(aga^{-1}) \geq \bar{\nu}(g)$, meaning that $\bar{\nu}$ is a normal \mathbb{L} -valued subgroup of G .

(e) Let $(G, \cdot, \mathfrak{N}) \rightarrow (H, \cdot, \mathfrak{M})$ be a continuous group homomorphism, and $\mu \in \mathbb{L}(H)$. Then

$$\begin{aligned} \overline{f^{\leftarrow}(\mu)}(e) &= \bigvee \{ \mathcal{F}(f^{\leftarrow}(\mu)) : \mathcal{F} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{F} \geq \mathfrak{N}^e \} \\ &\geq \bigvee \{ [e](f^{\leftarrow}(\mu)) : [e] \in \mathcal{F}_{\mathbb{L}}^s(G), [e] \geq \mathfrak{N}^e \} \\ &\geq \bigvee \{ \mu(e) : [e] \in \mathcal{F}_{\mathbb{L}}^s(G), [e] \geq \mathfrak{N}^e \} = \top, \text{ whence } \mu(e) = \top, \text{ since } \mu \in \mathbb{L}(G), \text{ implying } \\ &\overline{f^{\leftarrow}(\mu)}(e) = \top. \end{aligned}$$

Now let $x, y \in G$ and $\mu \in \mathbb{L}^H$. Then in view of the Definition 1(GLM3), we have:

$$\begin{aligned} \overline{f^{\leftarrow}(\mu)}(x) * \overline{f^{\leftarrow}(\mu)}(y) &= \bigvee \{ \mathcal{F}(f^{\leftarrow}(\mu)) : \mathcal{F} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{F} \geq \mathfrak{N}^x \} * \bigvee \{ \mathcal{G}(f^{\leftarrow}(\mu)) : \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{G} \geq \mathfrak{N}^y \} \\ &= \bigvee \{ \mathcal{F}(f^{\leftarrow}(\mu)) * \mathcal{G}(f^{\leftarrow}(\mu)) : \mathcal{F}, \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{F} \geq \mathfrak{N}^x, \mathcal{G} \geq \mathfrak{N}^y \} \\ &\leq \bigvee \{ \mathcal{F} \odot \mathcal{G}(f^{\leftarrow}(\mu)) : \mathcal{F} \odot \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{F} \odot \mathcal{G} \geq \mathfrak{N}_x \odot \mathfrak{N}^y \} \text{ (By applying Theorem 1.2.8 and Theorem 1.2.11[24], whence } f^{\leftarrow}(\mu) \in \mathbb{L}(G)) \\ &\leq \bigvee \{ \mathcal{F} \odot \mathcal{G}(f^{\leftarrow}(\mu)) : \mathcal{F} \odot \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathfrak{N}_{xy} \leq \mathcal{F} \odot \mathcal{G} \} \text{ (since } (G, \cdot, \mathfrak{N}) \in |\mathbf{S}\mathbb{L}\text{-NS}|, \text{ applying the Definition 10(LNGM), and due to Lemma 1, } \mathcal{F} \odot \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(G)) \\ &= \bigvee \{ \mathcal{H}(f^{\leftarrow}(\mu)) : \mathcal{H} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathfrak{N}_{xy} \leq \mathcal{H} \} = \overline{f^{\leftarrow}(\mu)}(xy). \end{aligned}$$

Finally, since $\overline{f^{\leftarrow}(\mu)} \geq f^{\leftarrow}(\mu)$, we get $\overline{f^{\leftarrow}(\mu)}(x^{-1}) \geq \overline{f^{\leftarrow}(\mu)}(x)$, for any $x \in G$. In fact, for any $\mu \in \mathbb{L}(H)$, $f^{\leftarrow}(\mu) \in \mathbb{L}(G)$ by Theorem 1.2.11[24]. So, we have:

$$\begin{aligned} \overline{f^{\leftarrow}(\mu)}(x) &= \bigvee \{ \mathcal{F}(f^{\leftarrow}(\mu)) : \mathcal{F} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{F} \geq \mathfrak{N}^x \} \\ &\leq \bigvee \{ \mathcal{F}^{-1}(f^{\leftarrow}(\mu)) : \mathcal{F}^{-1} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{F}^{-1} \geq (\mathfrak{N}^x)^{-1} \} \\ &\leq \bigvee \{ \mathcal{F}^{-1}(f^{\leftarrow}(\mu)) : \mathcal{F}^{-1} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{F}^{-1} \geq \mathfrak{N}^{x^{-1}} \} \text{ (by Definition 10(LNGI))} \\ &= \bigvee \{ \mathcal{G}(f^{\leftarrow}(\mu)) : \mathcal{G} \in \mathcal{F}_{\mathbb{L}}^s(G), \mathcal{G} \geq \mathfrak{N}^{x^{-1}} \} \\ &= \overline{f^{\leftarrow}(\mu)}(x^{-1}). \end{aligned}$$

Lemma 9. [2] Let $(G, \cdot, \Delta) \in |\mathbf{S}\mathbb{L}\text{-TOPGRP}|$, $\mu \in \Delta$ and $\nu \in \mathbb{L}^G$. Then $\mu \cdot \nu \in \Delta$.

Proof. Let $x \in G$, $\mu \in \Delta$ and $\nu \in \mathbb{L}^G$. Then $\mu \cdot \nu(x) = \bigvee_{st=x} \mu(x) * \nu(t) = \bigvee_{t \in G} \mu(xt^{-1}) * \nu(t) = \bigvee_{t \in G} \mathcal{R}_t(\mu)(x) * \nu(t)$. Fix $t \in G$, then $\nu(t)$ is constant and

$\nu(t) \in \mathbb{L}$. Since and $\mathcal{R}_t: G \rightarrow G$ is a homeomorphism, and $\mu \in \Delta$, $\bigvee_{t \in G} \mathcal{R}_t(\mu) \in \Delta$ and since Δ is stratified, and $(\mathbb{L}, *)$ is commutative semigroup, we have $\bigvee_{t \in G} \mathcal{R}_t(\mu) * \nu(t) = \nu(t) * \bigvee_{t \in G} \mathcal{L}_t(\mu) \in \Delta$, i.e., $\mu \cdot \nu \in \Delta$.

Proposition 3. [18] *Let $(X, \Delta_{\mathfrak{N}})$ be a stratified \mathbb{L} -valued topological space with a corresponding stratified \mathbb{L} -valued neighborhood system \mathfrak{N} . Then $(X, \Delta_{\mathfrak{N}})$ is Hausdorff-separated if and only if for all $x \neq y \in X$ there are $\nu_1, \nu_2 \in \Delta_{\mathfrak{N}}$ such that $\nu_1 * \nu_2 = \top_{\emptyset}$ and $\nu_1(x) * \nu_2(y) \neq \perp$.*

Definition 18. [18] *Let (X, Δ) be a stratified \mathbb{L} -valued topological space, $\mathfrak{N} = (\mathfrak{N}_x)_{x \in X}$ be the corresponding \mathbb{L} -valued neighborhood system, and A be a subset of X . Then closure of A , written as \overline{A} , is given by*

$$\overline{A} = \{x \in X : \mathfrak{N}_x(\top_{X \cap A^c}) = \perp\}$$

A subset of X is said to be closed with respect to Δ if $A = \overline{A}$.

Lemma 10. *A stratified \mathbb{L} -valued topological group $(G, \cdot, \Delta_{\mathfrak{N}})$ is Hausdorff-separated if and only if some singleton $\{a\} \subseteq G$ is closed. In particular $\{e\}$ is a closed subgroup of G .*

Proof. Let $\{a\} \subseteq G$ be closed subset of G . Then since the mapping $\varphi: (G \times G, \Delta \times \Delta) \rightarrow (G, \Delta), (g, h) \mapsto g^{-1}ha$ is continuous, we have $\varphi^{-1}(\{a\}) = \{(g, g) : g \in G\} \subseteq G \times G$, the diagonal which in view of the Corollary 6.2.1.2 [18], is a closed subset of $G \times G$ with respect to the product stratified \mathbb{L} -topology $\Delta \times \Delta$ implying that (G, \cdot, Δ) is Hausdorff-separated. Conversely, let $x \notin \{a\}$. Then $x \neq a \in X$ yields that there are $\nu_1, \nu_2 \in \Delta$ such that $\nu_1 * \nu_2 \leq \top_{X \cap \{a\}^c}$ and $\mathfrak{N}_x(\nu_1) * \mathfrak{N}_a(\nu_2) \neq \perp$, which implies that $x \notin \overline{\{a\}}$.

Lemma 11. *If $(G, \cdot, \Delta_{\mathfrak{N}})$ is a Hausdorff-separated stratified \mathbb{L} -valued topological group, and A be a closed subgroup of G , then the normalizer of A in $G: \mathbf{N}_G(A) = \{g \in G : \gamma_a(A) = A\}$ is a closed subgroup of G , where $\gamma_a: G \rightarrow G$ defined by $\gamma_a(g) = ag^{-1}a$ the conjugation map.*

Proof. If $a \in A$, take $c_a(g) = gag^{-1}$. Then the mapping $c_a: G \rightarrow G$ is continuous and hence the inverse image of the closed set $A: c_a^{-1}(A) = \{g \in G : gag^{-1} \in A\}$ is closed. Thus, we have

$$B := \bigwedge_{a \in A} c_a^{-1}(A) = \{g \in G : \gamma_a(A) \subseteq A\}$$

is a closed subset of G . Since the inversion mapping $j: G \rightarrow G, g \mapsto g^{-1}$ is a homeomorphism, A^{-1} is closed, since A is closed, and hence $\mathbf{N}_G(A) = B \cap A^{-1}$ is closed.

Lemma 12. *Let (G, \cdot, Δ) be a stratified \mathbb{L} -valued topological group, \mathfrak{N} be a corresponding stratified \mathbb{L} -valued neighborhood system on G and A is a subset of G . Then the centralizer*

$$\mathbf{Z}_G(A) = \{g \in G : [g, a] = e \quad \forall a \in A\}$$

is closed with respect to Δ . In particular, the center of G is closed subgroup.

Proof. If $a \in A$, then the mapping $\varphi : G \rightarrow G, g \mapsto [g, a] = gag^{-1}a^{-1}$ is continuous, where the element of the type $gag^{-1}a^{-1}$ is called *commutator* of the group G . Now since $\{e\}$ is closed subset of G , and since the inverse image of closed subsets under continuous mapping are again closed, in view of the Corollary 6.2.1.2 [18], $Z_G(a) = \{g \in G : [g, a] = e\}$ is closed, and as the $Z_G(A) = \bigwedge_{a \in A} Z_G(a)$ is closed, hence the result follows.

6. Conclusion

In this article, as a continuation of our previous work on \mathbb{L} -valued topological groups, where the underlying lattice \mathbb{L} was an enriched cl -premonoid, we have presented two types of results, one is about the relationship between \mathbb{L} -valued topological groups and their corresponding Kent convergence groups and conversely; the other is about \mathbb{L} -valued closure of \mathbb{L} -valued subgroup of a group. Although, it is an well-known fact that there is a close connection between principal limit convergence spaces and closure spaces, but we did not touch upon this issue here even for \mathbb{L} -valued generalization of these structures in conjunction with group structures, that is, to study \mathbb{L} -valued principal convergence spaces and \mathbb{L} -valued closure spaces. We intend to look into this issue in a future paper.

7. Acknowledgements

We are sincerely grateful to anonymous referees for generously giving their time to read our earlier version of this manuscript and providing various useful suggestions.

References

- [1] J. Adámek, H. Herrlich and G. E. Strecker. *Abstract and Concrete Categories*. J. Wiley & Sons, New York, 1990.
- [2] T. M. G. Ahsanullah. On fuzzy neighborhood groups. *J. Math. Anal. Appl.*, 130: 237–251, 1988.
- [3] T. M. G. Ahsanullah, D. Gauld, J. Al-Mufarrij and F. Al-Thukair. Enriched lattice-valued topological groups. *New Math. & Nat. Comput.*, 10(1): 27–53, 2014.
- [4] M. Baran. Closure operators in convergence spaces. *Acta Math. Hungarica*, 87(1-2): 33–45, 2000.
- [5] M. Baran, S. Kula, T. M. Baran, M. Qasim. Closure operators in semiuniform convergence spaces. *Filomat*, 30(1): 131–140, 2016.
- [6] M. Baran. Compactness, perfectness, separation, minimality and closedness with respect to closure operators. *Appl. Categor. Struct.*, 10: 403–415, 2002.
- [7] M. Baran. The notion of closedness in topological categories. *Comment. Math. Univ. Carol.*, 34(2):383–395, 1993.

- [8] R. Bělohlávek. *Fuzzy Relational System: Foundations and Principles*. Kluwer Academic Publishers, Dordrecht, 2002.
- [9] G. Birkhoff. *Lattice Theory*. Amer. Math. Soc. Colloq. Publ., Vol. 25, Providence, RI, 1967.
- [10] E. Čech. *Topological Spaces*. Revised ed. by Z. Forlík and M. Katětov, Wiley & Son, London, 1996.
- [11] S. C. Cheng, J. N. Mordeson, and Yu Yandong. *Lectures Notes in Fuzzy Mathematics and Computer Science*, Center for Research in Fuzzy Mathematics and Computer Science. Creighton University, Omaha, Nebraska, USA, 1994.
- [12] M. Demirci. On the convergence structure of L -topological spaces and the continuity in L -topological spaces. *New Math. & Nat. Comput.*, 3(1): 1–25, 2007.
- [13] D. Dikranjan, E. Giuli and A. Tozzi. Topological categories and closure operators. *Quaest. Math.*, 11: 323–337, 1988.
- [14] J. A. Goguen. L -fuzzy sets. *J. Math. Anal. Appl.*, 18: 145–174, 1967.
- [15] J. A. Goguen. Concept representations in natural and artificial languages: axioms, extensions, and applications for fuzzy sets. *Int. J. Man-machine Studies*, 6: 513–561, 1974.
- [16] J. Gutiérrez García, I. Mardones Pérez, and M. H. Barton. The relationship between various filter notions on a GL -monoid. *J. Math. Anal. Appl.* 230: 291–302, 1999.
- [17] U. Höhle and A. P. Šosta. Axiomatic foundations of fixed basis fuzzy topology, Chap. 3 *Mathematics of Fuzzy Sets: Logic, Topology, and Measure Theory*, The Handbooks of Fuzzy Sets Series, eds. U. Höhle and S. E. Rodabaugh. Vol. 3, Kluwer Academic Publishers, Dordrecht, 1999, pp. 123–272.
- [18] U. Höhle. *Many Valued Topology and its Applications*. Kluwer Academic Publishers, Dordrecht, 2001.
- [19] G. Jäger and T. M. G. Ahsanullah. Characterization of transitivity in L -tolerance spaces by convergence and closure, Submitted, Preprint, 2021.
- [20] D. C. Kent. Convergence functions and their related topologies, *Fund. Math.* 54: 125–133, 1964.
- [21] Y. C. Kim. Initial L -fuzzy closure spaces, *Fuzzy Sets and Systems*, 133(3): 277–297, 2003.
- [22] T. Kubiak. Separation axiom: Extension of mappings and embedding of spaces, Chap. 6, *Mathematics of Fuzzy Sets: Logic, Topology, and Measure Theory*, The Handbooks of Fuzzy Sets Series, eds. U. Höhle and S. E. Rodabaugh, Vol. 3. Kluwer Academic Publishers, Dordrecht, 433–479, 1999.

- [23] J. N. Mordeson, K. R. Bhutani, and A. Rosenfeld. Fuzzy Group Theory, in: Fuzziness and Soft Computing, Springer, 2005.
- [24] J. N. Mordeson and D. S. Malik. *Fuzzy Commutative Algebra*. World Scientific, Singapore. 1998.
- [25] J. N. Mordeson, and P. Nair. *Fuzzy Mathematics*. Springer-Verlag, Berlin, 2001.
- [26] A. Di Nola and G. Gerla. Lattice-valued algebras. *Stochastica* **XI, 2-3**: 137–150, 1987.
- [27] G. Preuss. Semiuniform convergence convergence spaces. *Math. Japonica*, 41: 465–491, 1995.
- [28] G. Preuss. *Foundations of Topology: An Approach to Convenient Topology*. Kluwer Academic Publishers, Dordrecht. 2002.
- [29] A. Rosenfeld. Fuzzy groups. *J. Math. Anal. and Appl.* **35**(1971), 512–517.
- [30] K. I. Rosenthal. *Quantales and Their Applications*. Pitman Research Notes in Mathematics. Vol. 234 Longman, Burnt Mill, Harlow, 1990.
- [31] E. Schechter. *Handbook of Analysis and its Foundations*. Academic Press, First Edition. October 30, 1996.
- [32] L. N. Stout. The logic of unbalanced objects in a category with two closed structures. Chapter 3; S. E. Rodabaugh et al (eds.), *Applications of Category Theory to Fuzzy Subsets*. Kluwer Academic Publishers, Dordrecht. 73–105, 1992.
- [33] C. L. Walker. Categories of fuzzy sets. *Soft Computing*, 8: 299–304, 2004.